


American Vegetable Grower

MAY • 1956




How to Kill 20 Million
Weeds an Hour!

The Vegetable Areas of
America—Louisiana

Five Ways to Control
Insects and Diseases

Facts You Should Know About Air-blast Spraying

Biggest money's worth in Pickups for '56



NEW FORD F-100 8-FT. 1/2-TON PICKUP,
GVW 5,000 lbs. Choice of 133-h.p. Six or 167-h.p. V-8.

Most Power!

Ford's new 1956 line of Pickups offers you the most power and performance in Ford history. Only the Ford Pickups give you gas-saving Short Stroke design in two engine choices, V-8 and Six! Ford's got the only modern Short Stroke Six in the 1/2-ton field!

Most Capacity!

Ford's new 8-ft. box gives you more loadspace than any other 1/2-ton pickup—up to 19 cu. ft. more. Available on the 118-in. wheelbase. Extra cost is surprisingly low. The standard 6 1/2-ft. box, on the 110-in. wheelbase, is one of the roomiest in its field—gives you a full 45 cu. ft. of loadspace.

Most Comfort!

Ford's new *Driverized Cab* is the easiest of all cabs to get in and out of. Doors open a full 70°—as much as 25° wider than other trucks. Most restful ride in any truck is yours with full foam rubber, 5 inches in seat, 3 inches in seat back . . . offered with 13 other extras in *Custom Cab*, low added cost.

Most Safety!

Only Ford Trucks have new deep-center Lifeguard steering wheel. Helps protect driver from contact with steering column in case of accident. Only Ford Trucks have new Lifeguard door latches, to help guard against doors springing open in an accident. No extra cost! Ford seat belts available at low added cost.

To get the most
for your old truck, see
your Ford Dealer now!



FORD TRUCKS LAST LONGER!

h



p. V-8.

you the most
Only the
stroke design
got the
n field!

dspace
u. ft. more.
cost is
, on the
n its field

of all cabs
—as much
ul ride in
5 inches
with 13
st.

r Lifeguard
contact
Only Ford
to help
accident. No
y added cost.

ER!




NU
DI

FU

Ther
corn
and
prev
ent

MAY, 19



**Correct
NUTRITIONAL
DEFICIENCIES**

**Control
FUNGUS
DISEASES**

There's a superior TC product to correct most nutritional deficiencies and TRI-BASIC COPPER SULFATE to prevent and control certain persistent fungus diseases.



NU-IRON

(A Nutritional Iron)

Nu-Iron is extremely effective for correction of chlorosis resulting from iron deficiencies . . . Results are noticeable shortly after Nu-Iron is applied. Easy and simple to feed to the plant through the leaves and stems by a direct application in spray or dust form.

TRI-BASIC COPPER SULFATE

A chemically stable copper fungicide containing not less than 53% metallic copper . . . For spraying or dusting truck and citrus crops. Especially effective in controlling persistent fungus diseases. Prevent fungus diseases through application of Tri-Basic Copper Sulfate before fungus attacks.

NU-Z

(Nutritional Zinc)

Nu-Z contains 53% metallic zinc . . . Can be applied directly to the plant in spray or dust form . . . Stimulates plant growth and corrects zinc deficiencies.

NU-MANESE

(Manganous Oxide)

An extremely effective nutritional manganese product for correcting manganese deficiencies due to low manganese content of the soil . . . Applied in spray or dust form.

COP-O-ZINK

(Nutritional Copper-Zinc)

This nutritional compound contains 48% copper and 4% zinc . . . Can be applied directly to the plants in spray or dust form. For correcting copper and zinc deficiencies and for stimulating healthier plant growth.

NU-M

(Nutritional Manganese)

A nutritional manganese compound to be fed to the plants through direct application in spray or dust form . . . To correct manganese deficiencies and to stimulate healthier plant growth.

ES-MIN-EL

and CUSTOM MIXED MINERAL MIXTURES

The essential mineral elements . . . Contains Manganese, Copper, Iron, Zinc, Boron and Magnesium, all essential to healthy, productive soil. Fruits and vegetables rich in vitamins cannot grow in soil poor in minerals. For soil application, ES-MIN-EL in spray or dust form for direct application to the plants is also available . . . Contains nutritional Manganese, Zinc and Copper.

WE WILL CUSTOM MIX MINERAL MIXTURES TO YOUR OWN SPECIFICATIONS IN LARGE OR SMALL QUANTITIES.

For Information on These
Nutritional Products,
Write, Wire or Phone Us.



DUST MIXTURES

Tennessee's Nu-Z, Nu-Iron, Nu-M and Tri-Basic Copper Sulfate are especially suited for use in preparing nutritional and fungicidal spray and dust mixtures and for use in mixed fertilizers.

TENNESSEE



CORPORATION

617-629 Grant Building, Atlanta, Georgia

Entered as second-class matter at Post Office at Willoughby, Ohio, under the Act of March 3, 1879. Additional entry at Mount Morris, Ill.

MAY, 1956


3

Bes•Blo's "Driving Air"

COVERS 80 FT.

in

ONE PASS



1956 Row Crop

BES•BLO

Blasts Pests with Air Power!

BES•BLO RC is the modern, time-saving, and more economical method of increasing row crop yields through better pest control. Its "Driving Air" penetrates foliage with uniform coverage of concentrate or dilute spray in widths up to 80 ft. Adjustable deflectors fit the Bes•Blo's spray pattern to every crop and weather condition.

In 3 Sizes...one to fit your needs. 200 lb. dust hopper (optional).

*Ask Your Dealer for a Demonstration
or write for information.*

BESLER

CORPORATION

4053 HARLAN STREET • EMERYVILLE, OAKLAND 8, CALIFORNIA

WAREHOUSE: Lansing, Michigan

AMERICAN
VEGETABLE GROWER

REG. U. S. PAT. OFF.
(Commercial Vegetable Grower)

Vol. 4 May, 1956 No. 5

FEATURED IN THIS ISSUE

Cover photograph by H. Armstrong Roberts.

Facts You Should Know About Air-Blast Spraying	9
By J. D. Wilson	
The Vegetable Areas of America—Louisiana	10
By Julian C. Miller	
Five Different Ways to Control Insects and Diseases	12
By Eldon S. Banta	
How to Kill 20 Million Weeds an Hour!	13
By Norman J. Smith	
Okra... The Gumbo Plant	20
He Bags His Carrots . . . and Regains Market	22
By Charles L. Stratton	
How to "Bag" Carrot Sales	23
Soil Insecticides . . . Your Underground Partner!	24
By D. E. Greenwood and R. N. Hofmaster	

DEPARTMENTS

Letters to the Editor	6
State News	14
Know Your Vegetable Seeds	14
Calendar of Coming Meetings and Exhibits	19
Answering Your Questions	35
New for You	36
Book Review	37
Editorial Page	38

AMERICAN VEGETABLE GROWER

Published Monthly by
AMERICAN FRUIT GROWER PUBLISHING CO.
Willoughby, Ohio
E. G. K. MEISTER,
Publisher
RICHARD T. MEISTER, Editor
Editorial Staff
E. K. GOULD, Managing Editor
H. B. TUKEY ELDON S. BANTA
R. L. CAROLUS M. P. RASMUSSEN
GEORGE M. ROSS, Art Director
J. S. BENDER, Production Manager
EDWARD L. MEISTER, Director of Advertising

District Advertising Offices
NEW YORK, Martin-Snow Co., 140 E. 46th St.
Phone—Plaza 5-0878
CHICAGO, Peck and Billingslea, Inc., 185 N. Wabash. Phone—Dearborn 2-0292
SAN FRANCISCO, McDonald-Thompson, 425 Market St. Phone—Yukon 6-0647
LOS ANGELES, McDonald-Thompson, 3727 West 6th St. Phone—Dunkirk 7-5391
SEATTLE, McDonald-Thompson, 1008 Western Ave. Phone—Elliot 3747
HOUSTON, McDonald-Thompson, 3217 Montrose Blvd. Phone—Jackson 9-4711
DENVER, McDonald-Thompson, 222 Colorado National Bank Bldg. Phone—Keystone 4-4669

AMERICAN VEGETABLE GROWER is published by American Fruit Grower Publishing Co., Willoughby, Ohio. E. G. K. Meister, Publisher and Chairman of the Board; Edward L. Meister, President; Richard T. Meister, General Manager; Gilbert Meister, Vice-President. Subscription price \$1.00 per year in U.S. and possessions, to Canada and other foreign countries \$1.50. Single current copies 25c, copies over one year old 75c.

When changing your address, send us old as well as new; send the address label from your last copy; allow 5 weeks for the first copy to reach your new address.

Postmaster: Please send change of address "Form 3579" to AMERICAN VEGETABLE GROWER, Willoughby, Ohio.

AMERICAN VEGETABLE GROWER

OWER

No. 5

UE

g Roberts.

at

..... 9

lea—

.....10

ol

.....12

.....13

.....20

.....22

.....23

.....24

.....6

.....14

.....14

.....19

.....35

.....36

.....37

.....38

OWER

ING CO.

or

S. BANTA

ASMUSSEN

ctor

ger

vertising

46th St.

185 N.

72

625 Mar-

3727 West

stern Ave.

Montrass

orado Ne-

4-4669

ublished by

Willingby,

Chairman of

Richard

ister, Vice-

year in

er foreign

Sc, copies

ld as well

last copy;

your new

ess "Form

VER, Wil-

GROWER

FRUITS Apples

Woolly apple aphid
Bud moth
Green apple aphid
Rosy apple aphid
Mealybug
Mites such as:
European red mite
Clover mite
Willow mite
Spider mites
Clinging mite
Fruit curculio
Red-banded leaf roller
Forbes scale

Pears

Mites
Pear psylla
Mealybug
Clinging mite
Fruit curculio
Fruit tree leaf roller
Red-banded leaf roller

Peaches

Mites such as:
European red mite
Two-spotted mite
Official mite
Fruit curculio
Codling moth
Orange tortrix
European Lecanium
scale
Soft brown
Aphids

Avocados

Latan
Green
Omniv
Orange
Soft bro

Mango & Passion Fruit

Fruit flies

Cherries

Black cherry
Black cherry
Fruit tree leaf

FRUITS Cranberry

Dates

Plums & Prunes

Strawberries

Blueberries

VEGETABLES

Leafhoppers
Black-headed
fireworms

Spittlebug nymphs
Crankberry fruitworm

Leafhopper, California and Southwest

Spider mites

Mealybugs

Spider mites

Mealy plum aphid

Plum curculio and

Mealy plum aphid

Aphids

Spider mites

Cranberry fruit worm

Cherry fruit worm

Blueberry maggot

Spider mites

Aphids

Colorado russet mite

Aphid

bean beetle

er

es

abbage

er

Cauliflower

Aphids

Cucumber

Aphids

Squash

Spider mites

Melon

Leafhopper on melon

Eggplant

Aphids

Peas

Spider mites

Beans

Aphids

Potatoes

Aphids

Onions

Trips

Spinach

Aphids

Turnips

Aphids

Brussels Sprouts

Aphids

Cabbage

Mites

Carrots

Cabbage looper

Leeks

Leafhoppers

Asparagus

Aphids

Garlic

Spider mites

Celery

Aphids

Alfalfa

Potato leafhopper

Spider mites

Alfalfa weevil larvae

Spittlebug

Vetch bruchid

Grasshoppers

Clover

Aphids

Mites

Young grasshoppers

Leafhoppers

Tobacco

Aphids


Not in recent years has any insecticide appeared that is so *useful* and—at the same time—so *usable* as malathion.

No other insecticide offers such high insect-killing power and such a high degree of safety to the user.

Malathion is accepted by the USDA for control of more than 80 insects on over 45 crops. It is the preferred insecticide for fly control . . . kills poultry mites, too.

And note these other advantages: You can apply malathion as close as 72 hours before harvest on many crops. And malathion is compatible with most other insecticides and fungicides.

Malathion insecticides are available from well-known manufacturers as wettable powder, emulsifiable liquid and dust. Order from your regular supplier. Do it today!




FREE: 1956 edition of MALATHION GROWER'S GUIDE. Write for your copy.

CYANAMID


AMERICAN CYANAMID COMPANY
AGRICULTURAL CHEMICALS DIVISION
30 Rockefeller Plaza, New York 20, N. Y.

NIAGARA "NO FROST"

25 Years' Successful Experience



Terminal Warehouse



Vegetable Freezing



Fruit Pre-cooling and Storage

Always full capacity

★

*Never any defrosting
shutdown*

★

*Always clean Freezing
or Storage Rooms*

★

*Write for Niagara
Bulletin 105*

NIAGARA BLOWER COMPANY

Dept. F.V. 405 Lexington Ave., New York 17, N. Y.

District Engineers in Principal Cities of U. S. and Canada

LETTERS TO THE EDITOR

Green Onion Bulbs

Dear Editor:

Along with my subscription to AMERICAN VEGETABLE GROWER, I am sending one of our new cartons of green onion bulbs. This package is being put up by Dutch Valley Growers, Inc., of South Holland, a farmers' co-operative.

Green onion bulbs are a selection of the better size bulbs from the onion sets. They



are the easiest and the most successful way to grow green onions either for sale or for the home garden. They can be started in the basement and even grown there if you have some sunlight. They can be planted outside in the latter part of winter and covered until springs sets in.

Ideal planting conditions are: After the frost is out of the ground, set in good loose soil up to the tip (either cover or leave uncovered). The plot should be well drained and fed with a high nitrogen fertilizer. Any good garden analysis will do. Keep the soil loose and the weeds under control. They will be the first vegetable in your garden ready to eat.

South Holland, Ill.

Judd Dalenberg

Marketing Peppers

Dear Editor:

I think your magazine is excellent in direct coverage of the vegetable grower's problems and clear, concise information regarding the latest improvements in the field. So much for so little.

Could you furnish us with some names and addresses of brokers, individuals, or produce houses in Illinois that would be interested in handling an early sweet pepper crop from this area? We are interested in developing this section as the land, climate, adaptability of machinery, etc., lead us to believe that we can produce this crop at least two weeks earlier than southern Illinois.

Wynne, Ark.

Jack Hooper

We sent our reader a list of the following wholesale buyers, brokers, and co-operatives in Chicago: L. Gillarde & Sons So., 141 W. Jackson Blvd.; D. M. Jacobson Co., Gridley, Maxon & Co., Inc., M. Lapidus & Sons, Riley-McFarland Co., S. H. Becker Co., Mushroom Growers Assn. Sales Co., and American National Foods, Inc., all located in South Water Market.—Ed.

AMERICAN VEGETABLE GROWER

ERS
ITOR

on to AMERI-
n sending one
onion bulb.
up by Dutch
th Holland, a
lection of the
on sets. They



successful way
or sale or for
be started in
there if you
be planted
winter and

e: After the
n good loose
ver or leave
ld be well
gh nitrogen
analysis will
e weeds un-
rst vegetable

d Dalenberg

excellent in
ole grower's
ormation re-
ents in the

some names
ividuals, or
it would be
sweet pepper
interested in
nd, climate,
lead us to
his crop at
n southern

ack Hooper
he following
o-operatives
ns So., 141
cobson Co.,
M. Lapidus
S. H. Beck-
Assn. Sales
oods, Inc.,
cket.—Ed.

LE GROWER

POWER COMMAND

IT'S ALL HERE IN THE...

ADVANCED

FERGUSON
SYSTEM

40



Vegetable growers everywhere will benefit from *Power Command* . . . the integration of the advanced Ferguson System with 4-Way Work Control and other Ferguson *firsts* to provide complete and instant mastery over tractor and implement operation.

See how the one tractor that dared to be different years ago still leads the way with five new models, every one of them with the famous Ferguson System: the new "40"; three "Hi-40" models . . . and a better than ever Ferguson "35". Ferguson, Racine, Wisconsin.

Ferguson

- ★ **NEW POWER STEERING**
Factory installed option.
- ★ **HEAVIER, LONGER MID-MOUNT, TOO!**
New Mid-Mount Cultivators available.
- ★ **NEW, 12-VOLT ELECTRICAL SYSTEM**
Hotter spark for surer starts.
- ★ **FREE VACATION TRAVEL CONTEST**
Ferguson Dealers have the details.


ORTHOCIDE 50 Wettable

(contains captan)

WINS FIRST PLACE

in potato and tomato fungicide comparison tests

Look at these results from 134 comparisons to ORTHOCIDE on 12 varieties of potatoes in 15 states and 3 Canadian provinces*



ON ALL CHEMICALS, READ DIRECTIONS AND CAUTIONS BEFORE USE.
T.M.'S REG. U.S. PAT. OFF.: ORTHO, ORTHOCIDE.



CALIFORNIA SPRAY-CHEMICAL Corp.

Executive Offices: Richmond, Calif.; Washington, D. C.

District Offices: Milwaukie, Oregon; Sacramento, San Jose, Fresno, Whittier, California; Phoenix, Arizona; Caldwell, Idaho; Shreveport, La.; Memphis, Tenn.; Maumee, Ohio; Haddonfield, New Jersey; Medina, New York; Columbia, South Carolina; Orlando, Florida.

Branch offices throughout U. S. A.


See for yourself in the result chart at the left. ORTHOCIDE 50 Wettable proved superior to other fungicides tested.

When you use ORTHOCIDE (contains captan) it can be like "extra money" growing on your potato plants. To get that "extra money" *this crop year*—start an ORTHO program featuring ORTHOCIDE now.

Tomatoes, too!

Tests at Rutgers University show ORTHOCIDE 50 Wettable increased

yield nearly 3 tons per acre over copper 53; exactly 3 tons per acre over competitive fungicide #1; 4¼ tons per acre over competitive fungicide #2 and nearly 5 tons per acre over untreated plants.



Proof like this points the way clearly to greater profit from better yields of better quality tomatoes. Isn't that what you raise them for? Then use ORTHOCIDE!

AMERICAN VEGETABLE GROWER

Facts You Should Know About AIR-BLAST SPRAYING

This new method of precision spraying means a big savings in water. Here are recommendations for getting the best results with air-blast equipment

By J. D. WILSON

Ohio Agricultural Experiment Station

WATER is the cheapest ingredient in a spray mixture—but vegetable growers are greatly interested in the possibility of using less of it in their spraying operations.

Water is sometimes in short supply at the spray site and may have to be hauled a considerable distance. This may require an extra piece of equipment. Besides this, it is in short supply in many parts of the country.



Concentrated sprays applied with air-blast equipment gave good control of diseases in tomatoes in Ohio experiments.

For these reasons air-blast spraying, which uses less water than commonly used methods of spraying for insect and disease control, holds a great appeal for many large growers.

So-called "concentrate" spray formulations may be applied with the commonly used hydraulic sprayer or with an air-blast machine. Air-blast sprayers now being used on row crops are designed to apply the spray material in a more concentrated formulation than the older, more familiar

recommendation of 2 to 4 pounds of the fungicide in each 100 gallons of water, applied at 125 to 200 gallons per acre.

Concentrated 4X and 5X formulations applied at 40 to 30 gallons per acre, respectively, are most commonly used in the air-blast spraying. The spray material is most often introduced into the air streams at 400 pounds per square inch at air velocities of 80 to 100 miles per hour.

Pump capacities of 15 to 20 and



The spraying job is done quickly and effectively with air-blast equipment. What the maximum spray swath should be is discussed in the article.

even more gallons per minute are most commonly used with the air-blast sprayers of today. However, many details of construction, design, and operation still remain to be determined in actual field operations before air-blast spraying obtains its maximum performance in the control of vegetable diseases.

Anyone using an air-blast sprayer for the first time should remember that the use of less water does not mean that the amount of fungicide applied per acre can also be reduced. For instance a 4X formulation would be required if the application of 40 gallons of the concentrate per acre were to correspond to the use of 160 gallons of an "X" formulation.

Also, it is poor practice to operate an air-blast sprayer in gusty winds or those in excess of 6 or 7 miles per hour. Furthermore, excessive swath widths should be avoided or the spray deposit in the outer limits (center of a double swath) may be insufficient to insure good disease control.

Performance studies with a variety

(Continued on page 32)

THE VEGETABLE AREAS OF AMERICA

LOUISIANA

By JULIAN C. MILLER

Louisiana State University

This is the sixth of a series on the important vegetable areas in the United States. New Jersey, Florida, Eastern Virginia, Arizona and Mississippi were covered in past articles.—Ed.

FROM its earliest settlement by the French, Louisiana has been known for its production of vegetable crops.

In the early 1800's river boats coming from Pittsburgh and St. Louis found that they could bring certain products, such as apples and potatoes, and on the return trip take back certain items, such as early potatoes, oranges and pecans. However, it was not until the Illinois Central Railroad was built from

However, as the population moved West many of the crop production areas moved with them. Carrots and onions moved to south Texas, and lettuce, garlic and artichokes to the West Coast. While they are still planted in south Louisiana, they are not produced in quantity as in times past.

Mixed Vegetables: For a long period of time Louisiana has been



LaSoda Irish potato developed at Louisiana station is high-yielding, early, red potato. Sailed not only to Louisiana and Gulf Coast, it is a leading variety in north central potato states.



Canfreezer pole bean, developed at the Louisiana station, is the leading commercial variety grown in Louisiana. It is ideal for processing.

Chicago to New Orleans that the vegetable area took on major proportions. This development took place just before the War Between the States and shortly thereafter.

Since these early periods south Louisiana has been a major section in the production of vegetables for shipment, as well as for local use.

noted for its shipment of mixed vegetables. These usually consist of shallots, carrots, and various greens such as mustard, turnips, beets, endive, and escarole, with shallots making the bulk of the load.

Louisiana is one of the few states that produce shallots in quantity today. This member of the onion family can be found on the market from November to May. The Louisiana station has bred varieties for all occasions, and one in particular, Evergreen, which does not die down or form sets during the summer. Therefore, the shallot can be found on the French Market at New Orleans practically every day during the year.


Green Beans: Both bush and pole beans make up an important crop

for fresh shipment and the third important crop for canning and freezing. They are grown in Plaquemines Parish (a county is called a parish in Louisiana), the Hammond area, and St. Martin Parish. Pole beans are grown largely in Plaquemines Parish and to some extent in southern Tangipahoa Parish.

Lima Beans: This crop is grown principally in the New Orleans area for the local market and for home use in other sections. Production is on the increase, primarily for the frozen industry.

Sweet Peppers: Louisiana holds a leading position in the production of sweet pepper. These are grown largely along the Mississippi River in St. James and St. John Parishes, and in the parishes of Tangipahoa

and St. M
to irrigate
Hot Pe
been note
ments.
in the Ne
Lafayette
years has
tion in
Gayenne,
principal
MAY, 1956




Commercial Vegetable Areas:

- 1—Sweetpotatoes
- 2—Irish potatoes
- 3—Beans (bush and pole)
- 4—Shallots and onions
- 5—Sweet pepper
- 6—Hot pepper
- 7—Okra
- 8—Southern peas
- 9—Watermelon and cantaloupe
- 10—Cucumber and squash
- 11—Cabbage, cauliflower, broccoli
- 12—Tomato
- 13—Mixed vegetables (mostly turnips, mustard, beets, carrots, collard)

grown commercially in the southern half of the state.

Southern Peas: This makes up another significant crop for the local market, shipment, and for freezing and canning. This crop is grown chiefly in the northern parishes, such as Jackson and Lincoln. How-



Unit 1 Porto Rico variety of sweetpotato developed by the Louisiana station comprises 80% of commercial crop grown in South. Photo shows ideal shape for Louisiana sweetpotatoes.

the third
ing and
n Plaque-
called a
ammond
sh. Pole
Plaque-
extent in
h.
is grown
ans area
or home
uction is
for the
holds a
oduction
e grown
pi River
Parishes,
agipahoa



Louisiana is one of only a few states that grow shallots in quantity. This field is the Louisiana Pearl variety of shallot.

Trellis on which sweetpotato vines have been trained in order to make crosses between varieties of the Louisiana University.



ever, it is grown to some extent in practically every section of the state. Southern peas are particularly prized as a southern food crop during the summer months.

Watermelons and Cantaloupes: These are grown in the sandy pine-land section of the state, principally in Ouachita, Jackson, Lincoln, Bien-ville and Washington Parishes.

Cucumbers and Squash: Cucum-bers and squash are shipped pri-marily from Tangipahoa Parish and sections in and around New Orleans. The fall crop is grown principally for shipment.

Cabbage, Cauliflower and Broc-coli: The local market area for these crops is south of New Orleans and the shipping area for these crops is centered in St. James and St. Mar-

(Continued on page 31)

and St. Martin, where it is possible to irrigate the crop.

Hot Pepper: Louisiana has long been noted for its hot pepper condiments. This industry is centered in the New Iberia-St. Martinsville-Lafayette area, which for many years has been the center of production in the United States. The Gayenne, Sport and Tabasco are the principal types grown.

Okra: The gumbo plant produces higher yields in Louisiana than in almost any other state. It makes up Louisiana's second canning and freezing crop, and its popularity for processing is expanding. It is also grown to a large extent for the fresh market. Many of the large soup manufacturers grow okra here, where it is frozen and transported to the manufacturing center. It is



Al Gerhart, North Ridgeville, Ohio, dusts cauliflower with duster he built. Unit is powered by Ford Model A engine. One drive is direct to the wheels, another goes to a second transmission that operates the duster. In this way duster can be operated independently and at various speeds from ground motion. High clearance permits dusting of tall crops such as sweet corn. Rear bar holding dust hoses is adjustable vertically. Scrap iron makes up bulk of framework. Because it is light in weight the machine does not pack ground.

Five Different Ways to Control INSECTS and DISEASES

Five eastern vegetable growers tell how they control pests to produce top-quality yields

By **ELDON S. BANTA**

PROFIT-MINDED vegetable growers keep one eye on economy and the other on effectiveness when planning their pest control programs. But they don't cut corners carelessly, you can bet on that!

This was the big lesson we learned from a recent swing through the East in which we visited five topnotch vegetable growers to see for ourselves how they are solving their insect and disease problems.

The Henry Way

Walter D. Henry and Sons, market gardeners of Eden, N.Y., market a fine crop of vegetables from their 300 acres. Sweet corn is an important crop, and corn borer is the big pest. On early varieties and plantings, control begins about June 10 with a 5% DDT dust. Three or four applications follow at intervals of four or five days. Later varieties and plantings are dusted when they get about knee-high, when the borer eggs reach the black head stage. Applications are repeated at five-day intervals until three or four applications are on. This program has also held corn earworm in check.

Tomatoes on the Henry farms are sprayed rather than dusted, and the chief problems are diseases. Early

blight is controlled with Zerlate and late blight with a Bordeaux or a 7% copper spray. So far these materials have not given the Henrys satisfactory control of anthracnose.

The first spray is applied about June 15 with repeat applications every 10 days until five or six have been applied. The first two are Zerlate sprays, the third Bordeaux, and the fourth Zerlate.

If late blight is found in the field at the time of the fourth spray, Bordeaux is again used. The fifth application is Bordeaux. Any later sprays depend upon weather and disease incidence.


Cucumber beetles are controlled with a 1% parathion dust. First application goes on when the first cotyl-

edon leaves open, and is repeated every five to seven days until beetles and other insects are no longer a problem. Anthracnose is controlled fairly well by applications of a 7% Zerlate dust beginning when plants first start to vine. Leaf blight is controlled by including a 7% copper dust with the Zerlate and parathion. Spotted mildew, a problem for many years, still defies complete control. The Henrys are trying Mildex and it looks quite promising.


Aphids on peppers have been successfully controlled by dusting as soon as they appear with a 1% parathion dust. The important step is to dust at the first sign of aphids.

The Adams Way

In the Hudson Valley east of Poughkeepsie, N.Y. Ralph Adams and Sons
(Continued on page 28)



Cucumbers that have a good control program produce vigorous vines like these and heavy crops.



Precision spraying equipment is almost a necessity for effective onion weed control. Boom built by John Youngs, Glen Head, Long Island, N.Y., has two nozzles per row which direct spray onto soil.



Unsprayed row at left presents a real hand weeding problem. Chickweed-free rows at right on Arthur and John Youngs' farm at Glen Head, L.I., received two 2-quart applications of Chloro IPC.

HOW TO KILL 20 MILLION WEEDS AN HOUR!

The new chemical weed killers are safe, quick, cheap, and easy to use—but go slow and follow directions

By **NORMAN J. SMITH**

*Associate County Agricultural Agent,
Nassau County, New York*

JUST ten years ago, if you had told your neighbors that your 14-year-old son could kill 20 million weeds an hour in a growing crop for 50 cents (the cost of 1 pint of 2, 4-D), they would probably have thought you were crazy.

To do a bit of quick figuring, a 20-foot boom sprayer operating at 4 miles per hour will spray 10 acres of corn an hour. Fifty weeds per square foot (a mild infestation) multiplied by 400,000 square feet equals 20 million weeds.

Chemical weed killers kill weeds easier, faster, cheaper, with rare crop damage, if used correctly, than other methods usually employed. A weed killer which doesn't meet these specifications soon passes out of use.

Let's review the weed killers that we have, by crops.

Asparagus: C. M. U. at the rate of 2 pounds in 50 gallons of water per acre, before the cutting season, and 2 pounds immediately after the cutting season, kills most broad-leaved weeds and grasses.

Beans: Sinox PE and Dow Premerge, one gallon in 30 to 50 gallons of water a few days before the beans

emerge, gives excellent control of broad-leaved weeds, and fair control of grasses.

Beets: Sodium chloride (salt) at 400 pounds in 200 gallons of water per acre applied when the beets have three to five true leaves. Purslane and lamb's-quarters are not controlled. There is room for a more effective weed killer on beets.

Carrots, Parsley, Parsnip, Fennel, and Dill: Stoddard Solvent, at 60 to 80 gallons per acre, kills most all weeds except ragweed. This weed

killer probably sold more vegetable growers on chemical weed killing than any other chemical. However, lower-cost materials would be welcomed for these crops.

Celery: In seedbeds only—Stoddard Solvent, 60 to 80 gallons per acre. Celery with "true" leaves can be injured.


Onions: Using potassium cyanate to control the weeds in onions has been an uphill fight from the beginning. Questions which faced the
(Continued on page 34)



Author stands in unsprayed mustard strip in Madison County, N.Y., oat field sprayed with 1/4 pound of 2,4-D when oats were 4 inches tall.

MAY, 1956

13



- Illinois Growers Form Co-operative-for-Profit
- Onion Hybrids and New Sweet Corn Varieties "on Trial" in New York

Co-operative-for-Profit

ILLINOIS—Vegetable growers of north Cook County announced formation of the Arlington Valley Growers Co-operative-for-Profit organization, and will file application for incorporation.

The growers' organization plans to issue 1000 shares of common voting stock having \$100 par value. Fifty charter members represent in excess of 2000 acres of vegetable production having an estimated value of \$1,250,000.

This is the first fresh vegetable co-operative-for-profit organization in Illinois, states William F. Lomasney, marketing specialist, University of Illinois Agricultural

ALLERGIC VEGETABLES

Sweet corn roots plowed under are toxic to tomatoes, lettuce, and onions, experiments at the University of Connecticut indicate. Plants become stunted with sickly root systems. By the time the soil warms up, the allergy causative agent is dissipated. Late-planted vegetables, such as beans, melons, and fall cabbage, do all right. Unfortunately, the allergy does not affect parsnips, a weed that seems to thrive following sweet corn.—Mrs. W. E. Griffith.

tural Extension Service. It is following the pattern of marketing organization work in heavily populated areas to bring about more orderly marketing, and to improve the growers' position in a more heavily integrated market.

The consumer will also benefit, because of this co-operative effort by getting fresher vegetables with a higher vitamin content than was possible when each grower packed and sold his own produce. The cost of pre-cooling, grading, and packaging equipment is so great that small growers—and they are in the majority in this area—were unable to offer vegetables in the best possible condition.

During 1956 the Arlington Valley Growers will grow and handle sweet corn, squash, parsnips, carrots, peppers, pumpkins, curly parsley, red and green cabbage, cucumbers, onions and onion sets, green onions, red beets, and turnips.

Earlier this month many of the vegetable growers who produce contract tomatoes for Campbell Soup Co. agreed to be represented by the Cook County Truck Gardeners & Farmers Association, for purposes of contract negotiations with the firm. Campbell officials agreed to recognize the organization and will consult with it in drafting next year's tomato contracts.—Elmer J. Steil, Assistant Sec'y, Des Plaines.

Interested in Chips?

OHIO—One out of every eight bushels of potatoes goes into the making of chips.

In a pamphlet issued by the National Potato Chip Institute, Dr. Ora Smith, research director, spells out the varieties which produce best for chipping. He cites the chippers' need for a potato well ma-

C7 LETTUCE AT MONACO

Taking advantage of a golden opportunity to get added consumer attention, C7 lettuce, a product of California and Arizona growers, was one of the sponsors of the CBS on-the-spot short wave rebroadcast of the colorful wedding and marriage of Grace Kelly and Prince Rainier III. C7 lettuce was also provided for the wedding feast; thus it capitalized on that facet of the well-publicized wedding as well.

The "Spring Salad Spree," of which the rebroadcast was an important part this year, is staged annually by Lettuce, Inc., at the peak of the Phoenix, Ariz., C7 season. This year's "Spree" is pushing C7 lettuce packaged in transparent film.

tured, high in specific gravity, and capable of making a light-colored chip.

Potatoes intended for the chip processor should be planted first as high specific gravity is brought about by maturity.

Copies of the pamphlet, entitled "Don't Get Caught Without Your Plans," are available on request to the institute at 946 Hanna Bldg., Cleveland, Ohio.

Late Spring

WISCONSIN—A late spring means that planting time for the potato crop will be

later than the last few years. Outlook indicated above average precipitation in April.

The value of seed treatment for potatoes and many of the early seeded vegetables will be of extra value under the cooler soil temperatures and higher moisture conditions that are indicated.

Potato growers might seriously consider seed treatment with thiram dust or similar materials. The value of insecticides for treatment of corn, bean, onion, and cucumber seeds along with a fungicide treatment will be good insurance against maggot injury to the seeds or young seedling plants.

We are recommending the use of lindane or dieldrin plus arasan on sweet corn and vine crop seeds; dieldrin, aldrin, or heptachlor plus arasan on onion seed; and lindane or dieldrin plus Spargon on bean seed to protect against maggot and wireworm injury and "damping-off."

The 1956 annual Wisconsin Potato Tour takes place July 24-27. Highlights will be visits to potato fields, machinery displays, annual golf tournament, and a day at the Spooner Branch Experiment Station.—John A. Schoenemann, Ext. Hort., Madison.

(Continued on page 16)

Know Your... VEGETABLE SEEDS

By VICTOR R. BOSWELL
U. S. Department of Agriculture

OKRA

SEED of okra is a true seed that develops in the okra pod, a fruit called a capsule. As it matures this many-celled pod becomes very fibrous and woody and then splits open along the length of each of the several long cells that contain the seeds. Under magnification the seed coat pattern of okra is rather pretty. Okra is an annual plant that is related to cotton and to the ornamental Hibiscus.

Growers and consumers outside the South may be surprised at the extent to which okra is grown. No dependable figures are available on the crop grown for food, but figures on seed supplies may give a hint. Eight pounds of seed is enough to plant an acre. About 1,100 to 1,200 acres was grown for seed in 1953 and 1954, respectively. Production amounted to some 340,000 and 590,000 pounds, or average yields of 300 and 500 pounds.

For many years thousands of tons of okra has been grown annually in the South for packing in brine and ultimate use in canned soups and related products. Only limited amounts are shipped fresh, but it is being quick-frozen to an increasing extent. Okra is grown almost universally in the South in home gardens. A large share of the seed surely goes to the amateur trade.

Okra seed has a relatively high oil content and has been considered for culture as



a source of edible oil. In contrast to many oily seeds, okra seed retains its viability for a long time under a wide range of storage conditions and is one of the best-keeping vegetable seeds. It presents no difficult problem at any stage of its production, recovery, or use for planting purposes.

VS


rs. Outlook
ipitation in

for potatoes
and vegetables
the cooler soil
ature condi-

ly consider
t or similar
cticides for
and cucum-
le treatment
maggot in-
lling plants.
e of lindane
et corn and
n, or hepta-
seed; and
on on bean
t and wire-

Potato Tour
ghts will be
ry displays,
day at the
tion.—John
adison.
(6)

DS



to many
bility for
storage
keeping
difficult
duction,
purposes.

E GROWER



Polyethylene film can help Control costly mite infestation

Large sheets of film made from BAKELITE Brand Polyethylene serve as fumigant barriers in the control of mites in the high-producing strawberry district of Watsonville, Calif. Resistant to weather, these polyethylene sheets are spread over the strawberry beds to contain the fumigant and hold it in contact with the plants during the period of fumigation.

Unaffected by most chemicals, flexible, and light in weight, the polyethylene film provides easy handling and does not flatten the plants over which it is spread. The same type of polyethylene film can also be used for a very effective mulch cloth.

Learn more about how film made of BAKELITE Polyethylene can help you on the farm—and in the market place as packaging for your produce. Write Dept. NS-66.

DID YOU KNOW: Pipe made of BAKELITE Polyethylene provides a new, easy, low-cost way to the finest farm water system. Pipe is easy to install with simple tools.




"Visqueen" film used in fumigating strawberries on the Tom Nakase farm was made by **The Vis-king Corp.**, Terre Haute, Ind., and fabricated by **Neil MacLean Co., Inc.**, San Francisco, Calif.



BAKELITE COMPANY, A Division of Union Carbide and Carbon Corporation **UCC** 30 East 42nd Street, New York 17, N. Y.
The term BAKELITE and the Trefoil Symbol are registered trade-marks of UCC


MAY, 1956

15




Concentrate Sprayer: two-way discharge, 22,000 cubic feet of air per minute from each side, 60 MPH. High pressure, abrasive- and corrosive-resistant, 20 GPM Myers piston pump. Easy-to-reach, positive controls.

Myers Air-Blast



Blower Attachment: 29" vaneaxial fan powered by air-cooled, 4-cylinder gasoline engine delivers 16,500 cubic feet of air per minute at 90 MPH. Readily adapts any Myers dilute sprayer to an effective concentrate rig. All controls can be reached from the tractor seat.

Field-Crop Sprayers



Two-way spray pattern produced by Myers 45,000 CFM field-crop, air-blast sprayers covers 80-foot spray swath. Turbulent roll of uniform air pattern assures complete coverage quickly and efficiently.

Deliver More Air



The combination of Myers high-pressure spray pump and long-lasting ceramic discs, provides ideal droplet size for concentrate spraying. The high-volume, high-velocity air stream saturated with man-made fog coming from spray nozzles, displaces the air around plants; covers leaves, fruit and twigs with uniform droplets of spray material.

For Best Coverage

Write today for the name
of your **Myers**® Sprayer Dealer

THE F. E. MYERS & BRO. CO., 8805 ORANGE ST., ASHLAND, OHIO
IN CANADA: THE F. E. MYERS & BRO. CO. (Canada) LTD., Kitchener, Ont.

STATE NEWS

(Continued from page 14)

Vegetable Testing Program

NEW YORK—The vegetable variety testing program for 1956 has been set up to include tests both at Ithaca and in various counties. At Ithaca preliminary screening trials of most of the newer varieties will be conducted.

It is planned to have a series of sweet corn demonstration plots and a series of onion trials to test some of the newer hybrids. Agents in 12 counties have indicated an interest in the sweet corn trials which will include a dozen of the newer varieties that have looked good in earlier screening trials.

The onion trials will include eight of the newer onion hybrids and will be located in the muck areas of Prattsburg, Elba, Oswego, Canastota, and Orange County.

In addition there will be a cabbage trial in Cortland County.—*P. A. Minges, Dept. of Vegetable Crops, Cornell University, Ithaca.*

Trellis Tomato Growers Join Hands

VIRGINIA—Trellis tomato growers in the Cape Charles section have organized the Virginia Trellis Tomato Association, in an effort to improve and standardize marketing of their product.

The Comet variety, which is grown exclusively in this section of the eastern shore, is a smooth, well-shaped, good-shipping tomato and has found wide acceptance in the trade due to its long shelf life at retail level.

Due to the trellis method of growing, very little off grade is produced generally. In the trellis method two rows of pruned plants are tied to a central overhead wire. The tent-shaped row thus formed allows the fruit to mature under the foliage without sun damage.

George Tinkham Sr., who introduced the trellis tomato to Virginia, was elected president of the association. Gilbert Doty is the new vice-president, and Robert Wyatt, secretary-treasurer. The organization represents 100% of the acreage to be produced this year in the area.

Attention! Prospective Queens

NEW JERSEY—Time's running out for prospective Vegetable Queens! Entries in this year's statewide vegetable queen contest must be submitted by June 18, reports James B. Woodford, chairman of the contest committee. Filled-in entry blanks (obtainable from farm organizations) should be sent to Mr. Woodford, Box 121, River-ton. Final judging will be made July 12 in the New Washington Hotel, Mount Holly.

Candidates must be 17 to 23 years of age and daughters of New Jersey vegetable growers. Top prize: \$100 and a \$150 wardrobe.

The committee is making a special study of ways for the queen to represent the state's vegetable growers and processors at public appearances throughout the year. Among other queenly attributes, the successful candidates must have poise and speaking ability.

More Tomatoes!

CALIFORNIA—There are 138,000 acres of tomatoes under contract this year as compared with 115,000 acres last year, estimates Howard Wilson, California Tomato Growers Association manager. Wholesalers' stock of canned tomatoes are sold and retailers' shelves are expected to be empty by harvesttime.

Vernon Glidden, superintendent for the Campbell Soup Company plant at Sacramento.—(Continued on page 18)

AMERICAN VEGETABLE GROWER

S
14)

ble variety
been set up
and in vari-
inary screen-
er varieties

ies of sweet
a series of
e newer hy-
ave indicated
trials which
er varieties
er screening

eight of the
be located in
burg, Elba,
ge County.
cabbage trial
inges, Dept.
University,

Hands
growers in
re organized
Association,
standardize

s grown ex-
the eastern
l, good-ship-
e acceptance
shelf life at

of growing,
ed generally.
s of pruned
erhead wire.
rmed allows
foliage with-

introduced
was elected
Gilbert Doty
and Robert
ne organiza-
creage to be

Queens

ing out for
Entries in
queen con-
18, reports
of the con-
blanks (ob-
121, River-
e July 12 in
ount Holly.
years of age
y vegetable
\$150 ward-

pecial study
represent the
rocessors at
t the year.
es, the suc-
poise and

8,000 acres
his year as
t year, esti-
nia Tomato
Wholesale-
are sold and
to be empty

ent for the
t at Sacra-
(8)

LE GROWER



Chomp, chomp, chomp the bugs are marching...

It's Spring again. And Colorado potato beetles, grasshoppers, cabbage loopers, onion thrips and other vegetable foliage destroyers are on the march. You can stop these pests—quickly and economically—with long-lasting dieldrin.

Dieldrin gives long residual

action. Long after application, it continues to give effective kill of foliage pests. Dieldrin is easy to use as a dust or spray. And dieldrin is economical, too. Mere ounces per acre provide dependable control.

This season be sure that foliage pests do not ruin your crops; steal

your hard-earned profits. Start the season off with an effective control program by applying dieldrin early . . . on the *crops* specified, in the *amounts* specified, and at the *times* specified. Dieldrin is available under well-known brand names from your insecticide dealer. See him today!

dieldrin
SHELL CHEMICAL CORPORATION
AGRICULTURAL CHEMICAL SALES DIVISION
460 Park Avenue, New York 22, New York



New for '56 two new models with Stainless Steel Tanks

For Spraying Dilute, Semi-Concentrates or Concentrates
The 200 Gallon Stainless Streamliner




200 Gallon
Stainless Steel Tank

7 1/2 G.P.M. Diaphragm
Pump or
30 G.P.M. Roller
Pump

The new "Stainless Steel Streamliner" with a 200 gallon stainless steel tank will give years of trouble free service at LOW initial cost. "ONE MAN OPERATION", light in weight (Less than 2000 lbs.). This combination dilute, semi-concentrate and concentrate sprayer is acclaimed as the outstanding sprayer of 1956. Other features include dual purpose for both orchards and row crops, adjustable axle as to height and width, and a choice of two special pumps for either dilute or high concentrate application.

The 100 Gallon Combination Buffalo Turbine Sprayer and Duster



Light enough to be
handled on "Hilly"
ground with the
SMALLEST TRACTOR
or
A "JEEP"

Large enough for the big grower and small enough for the small grower. The "Buffalo Turbine" combination Sprayer and Duster comes with a stainless steel 100 gallon tank, skid or trailer mounted. Designed for orchard or row crop spraying or dusting.

SPRAYS dilute, semi-concentrates or concentrate mixtures
DUSTS micronized chemicals, regular dusts or pellet baits
DISTRIBUTES Pellet or Granular insecticides or fertilizers

BUFFALO TURBINE
AGRICULTURAL EQUIPMENT CO.
INC.
GOWANDA, NEW YORK

Send For New 1956
Brochure and Literature

A Few Choice Territories
Open. Dealer-Distributor
Inquiries Invited.

STATE NEWS

(Continued from page 16)

mento, declared that his firm contracted for only 7% more than normal, but that canners have each signed a little more than normal, thus accounting for the increased acreage.—Neale Leslie



Photo courtesy Palm Beach (Fla.) Post-Times
Florida cucumber growers Jake Hammock (left) and Louis Alexander (right) discuss a three-week-old cucumber planting with Assistant Palm Beach County Agent John H. Causey. Cukes are planted in single beds with sunflower or sugarcane serving as windbreaks between every third bed. Hammock and Alexander grow squash, tomatoes, and cucumbers in partnership operations near Lake Park.

Ornamentals Carry Virus

OREGON—Greenhouse research by J. A. Milbrath and Roy Young, Oregon plant pathologists, has shown that Daphne odora from 48 different collections contained either cucumbers mosaic virus, alfalfa mosaic virus, or both—proving that cucumbers and the fragrant and beautiful Daphne odora do not go hand in hand.

Daphnes serve as intermediate hosts in the same fashion that wild gooseberries and currants serve as hosts for the wheat stem rust.

Daphne isn't the only ornamental known to carry viruses that infect food crops. Dahlias carry a virus known to infect tomato plants, as will flowering tobacco. Serious losses in commercial bean and cucumber fields have occasionally been traced directly to infection from gladiolus fields.

The idea is, say the pathologists, to keep these "poison-to-one-another" plants apart.—Harold and Lillie Larsen.

Irrigation in the East

PENNSYLVANIA—In the East, where rainfall is relatively heavy, irrigation should be only an auxiliary method to supply moisture, rather than the mainstay. Dr. Philip Minges, vegetable crops specialist at Cornell University, told a recent meeting of the Pennsylvania Vegetable Growers Association.

He said best yields and quality are obtained when plants have adequate moisture throughout the growing season, but warned against applying too much moisture.

No two farms can be irrigated in exactly the same pattern, Minges said. Type of soil, depth of soil, rooting habit of crops, and stages of plant growth, all need to be considered. He emphasized that in drought periods irrigation is the only solution.

This statement got full assent from Joseph Klock, Easton sweet corn grower, who said that irrigation last year saved him from a repetition of an almost complete crop failure in 1954 because of drought.

Klock cautioned growers to make a careful study of their irrigation needs and of different irrigation systems before they invest in "man-made rainfall."—N. M. Eberly, Agr'l Editorial Dept., University Park.

CALENDAR MEETING

May 10—Oration Greenhouse
Wittmeyer, D.

June 14-16—cluding Malb
convention, S
P. O. Box 11

June 27-29—Society annu
South Dakot
Brookings.—

July 24-27—Field days:
Stevens Point
Langlade Co
A. Schoenem
son.

July 28-A
Vegetable W

Aug. 1-Sep
tion of Ame
visiting Eng
Holland, Ge
Tour conduc
lean Express
land 16, Ohio

Aug. 11—Field Day, U
Oebker, 208
Urbana.

Aug. 14-1
meeting and
Wooster.

Sept. 7-9—Variety tria
ment Station
Minges, Cor

Sept. 23-6th Annual
Fontainebleau
Cooper, Exe
N. Y.

Sept. 27-able Assoc
tainebleau,
Colonial Dr

Oct. 31-N
and annual
Association
Sec'y, Box

Nov. 1-2-annual con
Francisco,
Los Angeles

Nov. 28-3
America, 4
Mich.—Jos
17th and I
6, D.C.

Jan. 28-3-able Ass
Philadelphia
St. N.W.,

CUCUMBERS

MORE tie
in the
Referen
publish
Sons, b
seeds.

Also
guide t
ease co
persona
rence
Calif.,

CALENDAR OF COMING MEETINGS AND EXHIBITS

May 10—Ohio Agricultural Experiment Station Greenhouse Vegetable Day, Wooster.—E. O. Wittmeyer, Dept. of Hort., Columbus 10.

June 14-16—Idaho Shippers Association (including Malheur County, Ore.) annual summer convention, Sun Valley.—Edd Moore, Exec. Sec'y, P. O. Box 1100, Idaho Falls.

June 27-29—South Dakota State Horticultural Society annual meeting, in conjunction with South Dakota Federation of Garden Clubs, Brookings.—W. A. Simmons, Sec'y, Sioux Falls.

July 24-27—Annual Wisconsin Potato Tour. Field days: July 24, Waupaca area. July 25, Stevens Point area. July 26, Antigo area and Langlade County. July 27, Spooner area.—John A. Schoenemann, University of Wisconsin, Madison.

July 28-August 4—Eighth annual National Vegetable Week sponsored by the VGAA.

Aug. 1-Sept. 12—Vegetable Growers Association of America, second Grand Tour of Europe visiting England, Norway, Sweden, Denmark, Holland, Germany, Switzerland, and France. Tour conducted by Walter F. Pretzer, % American Express Company, 1425 Euclid Ave., Cleveland 15, Ohio.

Aug. 11—Illinois State Vegetable Growers Field Day, University of Illinois, Urbana.—N. F. Oebker, 208 Vegetable Crops Bldg., U. of Illinois, Urbana.

Aug. 14-15—Ohio Pesticide Institute summer meeting and tour, Wooster.—J. D. Wilson, Sec'y, Wooster.

Sept. 7-8—Cornell Variety Field Days. Sept. 7, Variety trials at Ithaca. Sept. 8, Geneva Experiment Station and Robson Seed Farms.—F. A. Minges, Cornell University, Ithaca.

Sept. 23-26—Produce Packaging Association, 6th Annual Conference & Exposition, Hotel Fontainebleau, Miami Beach, Fla.—Robert A. Cooper, Exec. Sec'y, 500 Fifth Ave., New York, N. Y.

Sept. 27-29—Florida Fresh Fruit and Vegetable Association annual meeting, Hotel Fontainebleau, Miami Beach.—Geo. Talbot, 4401 E. Colonial Dr., Orlando.

Oct. 31-Nov. 1—Wisconsin State Potato Show and annual meeting Wisconsin Potato Growers Association, Antigo.—Harold R. Simmons, Exec. Sec'y, Box 22, Antigo.

Nov. 1-2—Western Growers Association 31st annual convention, Sheraton-Palace Hotel, San Francisco, Calif.—Headquarters 606 S. Hill St., Los Angeles.

Nov. 28-30—Vegetable Growers Association of America, 48th annual convention, Grand Rapids, Mich.—Joseph S. Shelly, Sec'y, 528 Mills Bldg., 17th and Pennsylvania Ave. N.W., Washington 6, D.C.

Jan. 29-31, 1957—United Fresh Fruit and Vegetable Association, Benjamin Franklin Hotel, Philadelphia.—Association headquarters, 777 14th St. N.W., Washington 5, D.C.

CUCUMBER MANUAL

MORE than 50 cucumber varieties are described and pictured in the new edition of "Descriptive Reference Manual of Cucumbers," published by Lawrence Robinson & Sons, breeders and growers of vine seeds.

Also included in the manual is a guide to cucumber insect and disease control, and a page for your personal field notes. Write to Lawrence Robinson & Sons, Modesto, Calif., for your free copy.

MAY, 1956

You can install your own irrigation system with USS NATIONAL* PLASTIC PIPE



*TRADE-MARK


Quick . . . Easy . . . Portable . . . Flexible

● Installing irrigation lines and sprinkler systems with lightweight polyethylene USS NATIONAL Plastic Pipe can be a one-man job. You can carry it anywhere—a 200-foot coil of two-inch National Plastic Pipe weighs only 86 pounds. It's easy to cut, and the fittings are of the insert type, quickly secured with stainless steel clamps. No adhesive necessary.

You can run flexible National Plastic Pipe anywhere—around corners or over obstructions. And it retains its toughness and flexibility over its entire working range from minus 90°F to plus 120°F.

USS NATIONAL Plastic Pipe is resistant to attack by rust, corrosion, acids, alkalies and salts. It won't crack or break with rough handling, even at very low temperatures. And due to its smooth interior surface, it offers very little flow resistance to the liquids passing through it. USS NATIONAL Plastic Pipe is available in sizes from 1/2-inch to 6 inches in diameter, in a variety of wall thicknesses.

For further details, write to National Tube Division, United States Steel Corporation, 525 William Penn Place, Pittsburgh 30, Pa. Ask for Bulletin No. 29.



SEE The United States Steel Hour. It's a full-hour TV program presented every other week by United States Steel. Consult your local newspaper for time and station.

NATIONAL TUBE DIVISION, UNITED STATES STEEL CORPORATION, PITTSBURGH, PA.
UNITED STATES STEEL EXPORT COMPANY, NEW YORK



NATIONAL plastic PIPE

UNITED STATES STEEL

**Normal rainfall
is NOT enough!**



FOR TOP YIELDS, MANAGE MOISTURE with GORMAN-RUPP IRRIGATION PUMPS

The big question about rainfall is not whether the year's total is normal, but whether enough falls at the right time. This rarely happens.

Even with normal rainfall, sprinkler irrigation makes sense. It makes even more sense to make

sure your system is the best with a Gorman-Rupp Irrigation Pump, backed by the combined experience of more satisfied users than any other make.

Be practical about irrigation. Get an engineered plan. No obligation. Write Gorman-Rupp today.

26 MODELS in centrifugal pumps for every irrigation requirement. Pump-and-engine packaged units with automatic check valve with level control, shaft seal under suction, large drum-type fuel tank—operational safety controls standard. Engines: air-cooled, radiator, heat exchanger. Gasoline, LP, natural gas, kerosene and diesel. Also PTO driven pumps.



THE GORMAN-RUPP COMPANY
305 Bowman St., Mansfield, Ohio

GROW and MARKET Quality Crops

The emphasis today is on quality. These books will help you produce the kind of crops the market demands.

- **DESTRUCTIVE AND USEFUL INSECTS.** By Metcalf, Flint, and Metcalf. 1,071 pages—\$12.50

A completely revised and up-to-date edition of this famous book. Valuable descriptions and photographs of all vegetable insects.

- **THE HOW-TO BOOK ON STRAWBERRIES.** By Robin Wyld. 112 pages—\$1.50

Written for the layman and illustrated with cartoons showing how to plant, tend, and harvest strawberries.

- **VEGETABLE GROWING.** By James S. Shoemaker. 515 pages—\$6.00

Second edition discusses individual vegetables, time and depth of planting, harvesting, and marketing.

- **THE CARE AND FEEDING OF GARDEN PLANTS.** 184 pages—\$3.00

The first book of its kind to acquaint the home gardener with plant hunger signs as they reveal themselves in lawns, on trees, shrubs, fruits, vegetables, house plants. Especially prepared color plates make it easy to identify the deficiency symptoms.

- **HUNGER SIGNS IN CROPS.** 370 pages—\$4.50

A plant may show unmistakable signs of hunger—nutritional deficiency—if we can only recognize them. *Hunger Signs in Crops*, especially prepared for the farmer or gardener, is a companion book to *The Care and Feeding of Garden Plants*.

All books sent postpaid on receipt of check or money order
AMERICAN VEGETABLE GROWER Willoughby, Ohio



Okra plant on farm of Orien Horn, Searcy, Ark. Plants grow, flower, and fruit from June to September, are at peak the first two weeks of August. Pods are picked six days after blooming.

OKRA... The Gumbo Plant

OKRA, a native of Africa, came to America around 1748, but only in the last 30 years has it attained commercial importance. Now okra is grown throughout the South, and is a basic ingredient of southern Creole dishes such as gumbo soup.

One of the largest centers of production is around Cairo, Ga., where some 6000 acres are grown annually. The bulk of the crop goes to the Campbell Soup Company's processing plant, the remainder to fresh market. Around Searcy, Ark., the Birds Eye freezing plant has some 1100 acres under contract.

Okra thrives in warm, humid climates, preferably where temperatures go above 85° F. It responds to plenty of water, but will endure drought quite well. The plants also respond to heavy fertilization.

Applications of 1000 to 1500 pounds of 8-8-8 per acre are suggested by C. D. Spivey, okra specialist at Georgia's Attapulgus Field



Harvesting okra from the Orien Horn planting at Searcy, Ark. Pods are snapped from the plants by hand. Pickers wear gloves to protect hands from stinging resulting from contact with pods.

AMERICAN VEGETABLE GROWER

Station. I
in before
side-dress

Okra
Searcy, A
3-9-27 per
bring his
duction in
2 tons of
pounds o
acre. He
4 tons of
the year l
a cover c
His yield
county.

Planted

Okra i
Seed is s
3½ feet
pounds p
ton, or b
rows 5 f
spaced 20
shallow c

Disease
major pes
damping-o
weather is

Fusariu
in many
should be
or main
bruised a
for the fu

Nemato
knot spec
hazard in
If a field
the soil s
planting
tode kille
ing okra
resistant

Harvest

Harves
after plan
peak dur
August. C
ing is ess
yields. I
plants, g
greatly r
picked at
Pods are
into soup
bloom.

Varieti
production
processing
Emerald
Campbell
widely gr
tain their
ing than
ties, alth
mands th
Clemson
Long Po
Hastings

MAY, 1956

Station. Half the fertilizer is disked in before planting; the remainder is side-dressed about six weeks later.

Okra grower Orien Horn, of Searcy, Ark., applies 200 pounds of 3-9-27 per acre before planting. To bring his soil up to good okra production he supplemented this with 2 tons of ground limestone and 300 pounds of 20% superphosphate per acre. He also applied between 3 to 4 tons of poultry manure per acre the year before, after plowing under a cover crop of Kentucky 31 fescue. His yields have been the best in the county.

Planted in Spring

Okra is planted about April 15. Seed is sown an inch deep in rows 3½ feet apart at the rate of 4 to 5 pounds per acre, using a corn, cotton, or bean planter. Horn likes his rows 5 feet apart and his plants spaced 20 to 24 inches apart. Early, shallow cultivation is essential.

Diseases and nematodes are the major pests. In early seedling stages damping-off is often a problem if the weather is hot and humid.

Fusarium wilt is a major disease in many stages of growth. Care should be taken not to bruise stalks or main roots in cultivating, as bruised areas make places of entry for the fusarium fungus.

Nematodes, especially the root-knot species, make okra growing a hazard in many parts of the South. If a field is known to be infested, the soil should be fumigated before planting with one of the new nematode killers. The practice of growing okra in rotation with nematode-resistant crops is desirable.

Harvest 60 Days Later

Harvest begins about 60 days after planting, and the crop is at its peak during the first two weeks of August. Careful, consistent harvesting is essential to obtain the highest yields. If old pods are left on the plants, growth stops and yields are greatly reduced. Pods should be picked at least three times a week. Pods are most suitable for processing into soups and gumbo six days after bloom.


Varieties vary with the area of production and the market. For processing in south Georgia, the Emerald variety developed by the Campbell Soup Company is most widely grown. The round pods retain their shape better after processing than do the star-shaped varieties, although the fresh market demands the latter. Other varieties are Clemson Spineless, Dwarf Green Long Pod, Louisiana Green Velvet, Hastings Dwarf.

THE END.

MAY, 1956

NO MITES

mean
better crops!



kill
and
control
mites
with

ARAMITE

There's more money for you from better crops. A small per-acre investment in Aramite can make a big dollar difference. Year after year, field results nationwide prove the bigger and better yields you can expect with Aramite—the safest, best-known mite-killer today.

Aramite gives outstanding control of most mite species including Two-Spotted, Atlantic, and European Mites. Equally effective on melons, citrus, corn, tomatoes, apples, peaches, and ornamentals. It is compatible with most insect and fungus-killing chemicals, has longer residual activity and is non-hazardous to humans and animals.

For more than 19 fruit and vegetable non-fodder crops, including lima beans, the Miller Amendment (Public Law 518) has approved Aramite with a tolerance established of one part in a million, because its disappearance is so rapid.



United States Rubber

Naugatuck Chemical Division

Naugatuck, Connecticut

producers of seed protectants, fungicides, miticides, insecticides, growth retardants, herbicides: Spergon, Phygon, Aramite, Synklor, MH, Alanap, Duraset.

LIFT OR LOWER — FLOOR TO FLOOR — WITH A STANDARD INCLINEBELT



Move boxes, cases, cartons, sacks or bundles from basement to first floor, or any floor to floor — continuously with the Standard Inclinebelt. Compact — simple to install — minimum maintenance and attention. Lifts or lowers 10 to 20 lbs. of live load per ft.; floor elevations of 8 ft. to 14 ft. 6 inches inclusive; two belt widths to handle commodities 15½ inches to 25½ inches wide. Electric motor operated. Write for INCLINEBELT Bulletin — address Dept. S-5.

STANDARD CONVEYOR COMPANY

General Offices: North St. Paul 9, Minn.
Sales and Service in Principal Cities



FERTI- LIQUID

Trade Mark Reg.

Patent Pending

The All-Purpose Liquid Fertilizer
Complete with Penetrating Agent —
Trace Elements — Growth Stimulant

**New, Advanced
10-20-10 Formula—
40% Nutrients**

Produces TOP QUALITY vegetable —BIGGER, BETTER-TASTING

- Gets it to market sooner
- BIGGER YIELD per acre
- Non-corrosive — use safely with any type spray (tractor, jet, boom or aerial)...ties in with insecticidal, fungicidal, herbicidal spray programs for low cost application
- No mixing, dust or odor!

\$1.85 per gal. in 55-gal. drums
del'd

Write for complete details

CLOVER CHEMICAL CO.

P.O. Box 10865 Pittsburgh 36, Pa.

He BAGS His Carrots... and Regains Market

New England grower prepackages top-quality product under brand name

By CHARLES L. STRATTON

CARROT grower Frank Palumbo, of Concord, Mass., has done a lot toward saving the New England market for New England growers.

Buyers haven't been too receptive to New England carrots due to the excellent quality of carrots from the West and the uneven bunches and poor quality sometimes put up by local growers. Palumbo, who specializes in growing high-quality carrots and was one of the first carrot prepackers in

an acre or several acres. His land is a rather sandy bottom loam. Hard ground, he claims, will produce misshapen carrots.

The only sidedressing Palumbo uses is nitrate and that only if necessary. A practice which he recommends to all growers is constant soil testing. "If rain washes the soil away," he says, "a soil test will tell right away where you stand and what the soil needs for the crop."

He uses chemical weed control, applying Salvat No. 5 according to directions, when the carrots get their true leaves. He prefers cooler weather for application as he finds there is not so much danger to plants then.

His entire program is based on producing an exceptionally nice line of carrots. One of his secrets is to seed thickly in order to get a slender carrot. No thinning is done in his fields as he prefers carrots with a diameter of three-quarters of an inch and less.

His prepackaging shed is a neat building in the center of a carrot field. The washer and packaging layout was



Carrots fresh from the washer (rear) are ready for grading and packaging in polyethylene bags under Palumbo's well-known Blue Ribbon label.


the East, has revitalized the local carrot market.

Each year he seeds some 50 acres of carrots in order to grow a uniform supply of tender, young carrots which he markets under his own Blue Ribbon label from about July 20 until the first week in November.

He doesn't use winter rye for early carrots as it takes too long to deteriorate. He likes to use 5-10-10, a ton to the acre. The ground is plowed, harrowed, then a three-quarter ton application of 5-10-10 to the acre is harrowed in and the ground is plowed again. He likes the ground loose and doesn't harrow again. The soil is broken up into beds and he allows 9 to 12 inches between the rows. The final application of fertilizer is applied and mixed with the soil and a four-row Model 300 Planet Jr. is used in the seeding.

Palumbo plants both the Hutchinson and the Imperator varieties. He claims the Hutchinson makes a nice early and a late carrot and is not as susceptible to blight. He uses the Imperator as a midseason carrot.

Carrots are planted as early as possible in the spring with the last planting before July 20. He plants a field at a time, whether it is a fraction of



Blue Ribbon carrots — 25 bags to the crate — await shipment to broker. Boxes are loaded on request of the buyer to insure crispness.

designed by a local concern (Larchmont Farms Co., Lexington, Mass.). Carrots are dumped directly into the hopper of a mechanical conveyor and moved up into the washer where they are given a thorough washing with cold artesian well water. The washer sides are constructed so the wind does not blow the water.

The carrots are spotless when they come out of the washer onto the table where they are sized and packed in polyethylene bags bearing Palumbo's Blue Ribbon trade name and his name and address.

THE END.

AMERICAN VEGETABLE GROWER

FRE
CH

This comb
polyethylene
sues in s
sues.

Ho
C

THE
bagging
dustry w
now tha
marketin
ers and
the effec

Result
cate tha
helped s
ducted
iversity
markets
Pittsbu
from Ap

Method

The m
1) On
units of
2) On
with a t
than bun
3) A
and two
with the
cents les

4) A
pound p
pound bu

The la
from the
and two
Average
amounte
per 100
than one
the leas
pound bu
per 100 c

Sales a
100 custo
display o
bags and
pounds p
display o
bags.—W

MAY, 1956



This combination display of one- and two-pound polyethylene bags of carrots produced largest sales in study of bagged and bunched carrot sales.

How To "Bag" Carrot Sales

THE switch from bunched to bagged carrots hit the carrot industry with dramatic suddenness. But now that we are used to the idea of marketing carrots in bags, some growers and retailers are wondering what the effect has been on carrot sales.

Results of a sales experiment indicate that bagging carrots may have helped sales. This study was conducted by Pennsylvania State University and the USDA in 12 supermarkets of the Kroger Company in Pittsburgh during four test weeks from April 4 through May 7, 1955.

Methods of Display

The methods of display were:

- 1) One-pound bunches priced in units of two.
- 2) One-pound polyethylene bags with a two-unit price four cents less than bunches.
- 3) A combination display of one- and two-pound polyethylene bags with the two-pound bag priced two cents less than two one-pound bags.
- 4) A combination display of one-pound polyethylene bags and one-pound bunches priced as in 1 and 2.

The largest sales were obtained from the combination display of one- and two-pound polyethylene bags. Average sales from this display amounted to 7.3 pounds of carrots per 100 customers. This was more than one-third larger than sales from the least effective display of one-pound bunches which sold 5.6 pounds per 100 customers.

Sales amounted to 6.9 pounds per 100 customers from the combination display of one-pound polyethylene bags and one-pound bunches, and 6.6 pounds per 100 customers from the display of one-pound polyethylene bags.—W. A. Lee, Penn State U.



TAWCO Leaf Crop Harvester

For—

Spinach
Mustard
Greens
Kale
Salad Greens
Turnip Greens
Parsley

Combat rising labor costs in harvesting market greens with the Tawco Leaf Crop Harvester. Gas engine powered and single man operation. Cuts foliage above or below ground at desired height, and conveys it without bruising to chute where it drops into basket. After basket is filled, foliage accumulates in chute for hand compression to give desired pack in basket.

Works equally well on trench irrigated crops or flat planting. Growers producing for canning or freezing find Tawco Harvester recovers much higher percentage of usable foliage than larger, more costly machines. Any grower of size quickly pays for Tawco Leaf Crop Harvester out of labor and crop savings affected.

TAWCO Radish Harvester and Topper

Has won wide acceptance since introduction in 1954. Operates on similar principle to Leaf Crop Harvester except that root is recovered and tops are discarded. One large grower with nine machines harvests 15 acres a day. He replaced 160 field and packing shed employees with nine workers—paid for his machines from a few days' savings in labor costs.

For Prices and Descriptive Literature Write

TAWCO Products, Inc.

Vegetable Harvester Division

1224 Chesapeake Ave.

Columbus 12, Ohio

PROFITABLE ROADSIDE MARKETING

By R. B. Donaldson
and W. F. Johnstone


A practical handbook for the successful operation of a roadside market. It is the result of many years' study of "on-the-farm" marketing in all parts of the country by two experts in the field.

Every phase of roadside marketing is covered—planning, building roadside stands, handling and storing produce, packaging, pricing, advertising, etc. Illustrated with photographs, charts, drawings.

Sent postpaid on receipt of \$2.00

AMERICAN VEGETABLE GROWER
Willoughby, Ohio

Easy-to-Build Roadside Stand



This roadside market, conveniently located so the motorist can readily stop, will make it easy for you to take care of customers quickly. Movable racks make it possible to maintain an attractive stand in accordance with supply and demand. Overhanging roof affords protection for the customer.

Working drawings sent postpaid on receipt of \$1.50
AMERICAN VEGETABLE GROWER
Willoughby, Ohio

Now!... A PORTABLE Stericooler

**THAT PUTS
HYDROCOOLING
IN THE GROWERS'
BACKYARD
at the peak
of the harvest!**



Right side of Portable Unit with jacks in working position.

Here is a big capacity Stericooler hydrocooling unit designed and engineered to meet seasonal requirements of buyers and shippers of perishable fruits, vegetables, berries.

Follow the crops in season with your portable Stericooler hydrocooler—handle from 200 to 225 bushels of produce an hour—ship when the market is "hot" and take full advantage of the extra profits that Stericooler hydrocooled fruits and vegetables always bring.

The Stericooler hydrocooler is specially designed and engineered for mounting on a semi-trailer. It has two ample working platforms, four landing jacks for stability; plus rugged construction and engineered efficiency found only in FMC Stericooler hydrocooling units.

The portable Stericooler hydrocooler is available on a trailer ready to go. Or if you prefer we will supply the portable Stericooler unit ready for mounting on your trailer with complete mounting instructions.

Portable unit showing ice tank, pump, gas engine drives. (Also available with electrical motor drives.)



**GET YOUR SHARE OF STERICOLER
HYDROCOOLING PROFITS—
MAIL COUPON TODAY!**



Gentlemen: I'm interested in increasing my profits with your Portable Stericooler hydrocooler. Please send more information.

NAME _____
ADDRESS _____
RFD NO. _____ SC6A

FLORIDA DIVISION
FOOD MACHINERY AND CHEMICAL CORPORATION
P. O. BOX 1718 • LAKELAND, FLORIDA

SOIL INSECTICIDES ... Your Underground Partner!

Irish potato and sweetpotato growers have found a strong ally in the war against soil insects

By D. E. GREENWOOD and R. N. HOFMASTER
Virginia Truck Experiment Station

SOIL insecticides are effective primarily against wireworms, grubs, and similar insect forms which live exclusively in the soil and do most of their damage to underground portions of plants. In fact wireworms, more than any other single pest, were largely responsible for the development of modern soil insecticides because of the tremendous losses they inflicted on root and transplant crops at a time when effective and practical control methods did not exist.

Wireworms are now effectively controlled, in most soil types, by the application of from 2 to 3 pounds of aldrin or heptachlor per acre.

Zone of Activity

Wireworms are slender, hard-shelled, amber colored worms. Actually, they are the young or immature form of "click beetles." During the winter wireworms go down below the frost line; in the spring they move up to the surface foot of soil and remain there for greater part of the growing season. Once in this zone, their lateral movements (except for a certain amount of aimless wandering) are in response to a search for food.

Soil insecticides applied broadcast and worked into the top 5 or 6 inches of soil thus assure an almost continuous period of contact between the wireworms and the material used, whether the worms are actively feeding or not. As most of the vulnerable plant parts are also contained within this soil zone, control is usually



Flea beetles in untreated Irish potato plot at foreground have completely defoliated plants; treated plots at rear (aldrin) and right (heptachlor) have good foliage, show no damage from flea beetles.

achieved before the worms can inflict much damage.


Potatoes Down-graded

Irish potatoes and sweetpotatoes developing in untreated, infested soils are subject to intermittent periods of intensive wireworm feeding. While the familiar feeding holes, as shown in the roots of sweetpotato, do not greatly impair the quality of the roots or tubers, they down-grade them, with a proportionate loss to the grower.

In a recent test with Irish potatoes, in which aldrin or heptachlor had been applied broadcast at the rate of from 2 to 3 pounds per acre, the treated areas averaged less than .01% injury, while the untreated portions of the field averaged close to 30% injured tubers. This can be considered as a serious infestation because to bring such a field up to grade requirements would necessitate the most rigorous sorting table technique. The cost of the material alone would hardly pay one man's wages for one day at the grading table.

Another serious wireworm problem, though less conspicuous than potatoes full of holes, results from wireworm feeding on sprouting seeds. The seeds are generally hollowed out and the stand thinned, but occasionally the growing point is nipped on a seed that has already sprouted. When such plants come aboveground, they are malformed and useless.

(Continued on page 26)



At top, sweetpotato roots from untreated land infested with wireworms have numerous black feeding holes which down-grade product. Below, typical undamaged roots from plots treated with aldrin or heptachlor at 2 to 3 pounds per acre.

po
ag

Stau
for l

Everyo
formul
is easil
400 is
insecti
other i
bulleti

Stau

Damp-
tomato
halted
yields,
Stauffe
We hav

Stau

Kill th
you'll h
4-S do
you can
several
for a S
*formerly

STA
380 M

Stauffer
New Yor

MAY, 19

ner!



plot of fore-
ants; treated
achlor) have
lea beetles.

can inflict

etpotatoes
ested soils
periods of
g. While
as shown
o, do not
y of the
own-grade
loss to the

n potatoes,
chlor had
the rate of
acre, the
than .01%
d portions
e to 30%
be consid-
because
grade re-
e the most
ique. The
ne would
es for one

orm prob-
ous than
ults from
ting seeds.
ollowed out
occasion-
pped on a
ted. When
ound, they
s.

26)

LE GROWER

3 powerful allies against Insects and Disease

Stauffer's "FLOWABLE" PARATHION 400* for Insect Protection

Everyone likes the powerful insecticidal properties of Parathion, but here's a formulation which has no dust hazard... which has little tendency to splash and is easily washed from skin and clothing if it does. Stauffer's "flowable" Parathion 400 is an emulsion in water... broadly compatible with fungicides and other insecticides... low in phytotoxicity... high in wettability... without solvent or other inert residues to affect flavors or injure plants. Ask us to send a Stauffer bulletin on "Flowable" Parathion 400.

Stauffer's CAPTAN 50-W for Disease Protection

Damp-off and root rot on beans and peas, early and late blight on potatoes, tomato blights, and the numerous diseases that attack cucurbita and carrots are halted by Stauffer's CAPTAN 50-W. Our customers tell us they enjoy higher yields, bigger size, exceptional finish and color, unusual keeping qualities... when Stauffer Captan 50-W is used to control these characteristic diseases of vegetables. We have—and will gladly send—an informative bulletin on Stauffer's Captan 50-W.

Stauffer's VAPAM 4-S for Your Seed Beds

Kill those weeds, germinating weed seeds, nematodes, soil fungi and insects and you'll have a big growth of healthy plants to set out. One application of VAPAM 4-S does the work... no special equipment is needed... no ground coverings... you can treat acres in a day! VAPAM has been used with remarkable success in several states and it is now available nationally. Send for a Stauffer VAPAM Vegetable seedbed bulletin.

*formerly Stauffer's Paraflow.



STAUFFER CHEMICAL COMPANY
380 MADISON AVENUE, N. Y. 17, N. Y.

Stauffer products—"Everything for the Grower"—include Oxex, Dieldrin, Malathion, Ceramite, Systox, DDT, BHC, Sulfenone and Sulfurs. New York • San Francisco • Houston • Los Angeles • No. Portland • Lubbock • Harvey • Weslaco • No. Little Rock • Omaha • Apopka

"Flowable" Parathion Controls

Army worms
Bud moth
Cabbage worm
Catfacing insects
Codling moth
Corn borer
Cottony cushion scale
Cottony peach scale
Cucumber beetle
Florida red scale
Forbes scale
Fruit tree leaf roller
Grasshoppers
Green peach aphid
Green stink bugs
Leaf hopper
Leaf miner
Mealy bug
Mealy plum louse
Mexican bean beetle
Oriental fruit moth
Peach tree borer
Pear blister mite
Pear psylla
Plum curculio
Purple scale
Red-banded leaf roller
Red bug
Red spiders
Rosy apple aphid
San Jose scale
Scurfy scale
Spider mites
Thrips
Twig girdler
Webworms
Woolly apple aphid

Stauffer CAPTAN Controls

On apples: Scab, sooty blotch, fly speck, bitter rot
On carrots: Septoria, cercospora
On cherries: Leaf brown rot
On cucurbits: Angular leaf spot, anthracnose, downy mildew
On ornamentals: Damp-off
On peaches: Brown rot and peach scab
On potatoes: Early & late blight
On roses: Black spot
On strawberries: Leaf spot and grey mold fruit rot
On tomatoes: Early & late blight, anthracnose, stemphylium


Stauffer VAPAM Kills

Amaranthus sp.
Armilaria
Bermuda grass
Chickweed
Dandelion
Fusarium sp.
Johnson grass
Lambs' quarters
Malva
Nematodes
Nut grass
Phytophthora sp.
Purslane
Ragweed
Rhizoctonia sp.
Russian knapweed
Symphilitis
Verticillium
Water grass
Wild morning glory

STANDARD *ize*

on SINOX PE

Amine Dinitro Weed Killer
the pre-emergence
chemical weed control
for peas, beans and sweet corn



SINOX PE eliminates the weeds that are robbing your crops of valuable nutrients, without harming the crops themselves . . . does the work of a whole gang of field men at a fraction of the cost! This profitable spray can be applied at *any time* between planting and emergence . . . whenever it is most convenient for you.

Ask your Dealer for complete SINOX PE literature today, and consult local authorities for recommendations.

Remember — good weed control is good farm economics . . .

...it pays to
Standard-ize



**standard
agricultural
chemicals, inc.**

1301 JEFFERSON ST., HOBOKEN, N. J. • 429 FORUM BUILDING, SACRAMENTO, CAL.

SCOTT-URSCHEL Red Beet and Carrot Combine



**DIGS-
TOPS-
and LOADS**
in one
OPERATION

Here's an improved version of the famed Scott combine that gives you the operational features you want. Ideal for Cello-Pack carrots as well as for canning carrots and red beets. Topping is as close to the carrot or beet as possible, yet with no damage to the commodity. High speed of harvesting eliminates need for stock-piling . . . harvest as the canner or packer needs the crop. Write today for complete information on Model 004-R for red beets and carrots for canning, or Model 004-RX for fresh market carrots.

Scott Viner Company 1224 Kinnear Road Columbus 8, Ohio

SOIL INSECTICIDES

(Continued from page 24)

In years past, the only practical means of combating this problem was to continue planting until the wireworms had their fill. The lima bean industry on the West Coast for many years struggled along at great cost because of this type of wireworm injury. Sweet corn planted in infested fields is particularly vulnerable.

Earlier Yields

The value of soil insecticides was clearly demonstrated in conjunction with the potato experiment referred to above. One grower made several plantings of early cucumbers. One planting was made on soil which had been treated for wireworms; the other plantings were made on untreated, infested fields. The plantings on treated soil resulted in a good stand the first time over and was actually yielding pickling cukes at a time when the other fields were just getting established.

The advantages of getting a crop up and into production for those all-important first few days of a good market are obvious. This is particularly true of a high cash crop like pink and green-wrap tomatoes, an up-and-coming industry in Virginia which has run into the problem of wireworms entering the fruits which come in contact with the soil. In such special cases, soil treatment has become almost a routine practice.

Transplant crops such as tomatoes, peppers, and cabbage that are usually started in seedbeds and later transferred to the field, are particularly vulnerable to wireworm feeding, especially if the wireworms have not previously been able to obtain food from a cover crop. Wireworms enter the underground stem and bore up; causing the plants to wilt and die. In fields known to be infested, it is unwise to plant such crops without first having applied a soil insecticide.

Effect on Foliage Feeders

One of the more recent developments in the expanding research on soil insecticides has been their application to problems dealing with pests normally considered as foliage feeders such as the potato flea beetle. This pest can chew so many holes in potato foliage that the effective leaf area of the plant is reduced to practically nothing and all growth stops.

Since this injury comes after tubers have started to form, the plant foods required for tuber enlargement are not available. Reduction in yield is brought about by a loss in size of tubers, not in their numbers.

The potato flea beetle is a small, black insect about 1/16 inch in length.

AMERICAN VEGETABLE GROWER

In Virginia adult stag nation in a convenient ings or oth beetles dep soil at the

The you hatch from ground pl manner as to tubers occasion c worm inju

After d pleted in t adults em is the gen structive t fields can week. Aln ing in any



Charles V. L drew, of So loaded plant of Cobden, heavily to g which is hig cents a pound. Lamer set shorter solu Just after pl acre of 3-1/2 growing well plied per ac plant by han over by July

from eggs beetles ear

Tests h of either a applied to decrease beetles, th tuber dam yield over age applic ferred fol not signif

Another of soil ap cation lie timing. S the spring planted an out the em treatment affected b means th the dama

DES
(24)

practical
blem was
the wire-
ima bean
for many
reat cost
wireworm
infested
ble.

cides was
njunction
referred
several
ers. One
which had
ms; the
on un-
plantings
a good
was actu-
at a time
just get-

g a crop
those all-
f a good
particular-
like pink
a up-and-
ia which
of wire-
ich come
such spe-
s become

tomatoes,
e usually
er trans-
particularly
ding, es-
have not
ain food
ms enter
bore up;
d die. In
it is un-
out first
icide.

develop-
earch on
eir appli-
with pests
age feed-
ttle. This
in potato
f area of
practically
er tubers
nt foods
ment are
yield is
e of tub-

a small,
n length.
GROWER

In Virginia it overwinters in the adult stage and emerges from hibernation in April, just in time to move conveniently onto new potato plantings or other hosts. In early May, the beetles deposit tiny white eggs in the soil at the base of the host plant.

The young flea beetle larvae which hatch from these eggs feed on underground plant parts, much in the same manner as wireworms; in fact, injury to tubers by the flea beetle larvae on occasion can be just as severe as wireworm injury.

After development has been completed in the soil, a new generation of adults emerges in early June. This is the generation which can be so destructive to potato foliage and whole fields can be skeletonized within a week. Almost all of the beetles feeding in any given field have developed



EARLY TOMATOES

Charles V. Lamer (left) shows Dr. William Andrew, of Southern Illinois University, a heavily-loaded plant of Big Boy tomatoes on his farm at Cobden, Ill. Lamer fertilizes early tomatoes heavily to get yields of 10 pounds per plant, which is high for his area. He gets 10 to 20 cents a pound for them.


Lamer set his plants early in April with a starter solution to give them an early boost. Just after planting he applied 1200 pounds per acre of 3-12-12 fertilizer. When plants were growing well, 800 pounds of 3-12-12 were applied per acre, with applications made to each plant by hand. Harvest began in June and was over by July 25.—E. S. Banta.

from eggs laid by the overwintering beetles early in the spring.

Tests have shown that 2½ pounds of either aldrin or heptachlor per acre applied to the soil resulted in a 94% decrease in foliage injury by flea beetles, the practical elimination of tuber damage, and a 75% increase in yield over untreated fields. Four foliage applications of dieldrin, the preferred foliage recommendation, were not significantly better.

Another very important advantage of soil application over foliage application lies in the matter of proper timing. Soil applications are made in the spring before the potato crop is planted and remain effective throughout the entire growing season. Foliage treatments, on the other hand, can be affected by seasonal variation, which means they are often applied after the damage has been done. THE END.

MAY, 1956



Successful growers by thousands have shown that our baskets, boxes, and containers for retail and bulk marketing promote better profit for the vegetable producer.

Product of our complete PLANNED PACKAGING facilities, these marketing specialties are made of fine quality corrugated board, attractively printed in 2 colors. Sizes—2, 4, 8, 12, and 16 quarts. Wood or wire handles. Shipped flat for easy storage. Easy to set up. Write PRODUCE SALES, The Ohio Boxboard Co., Rittman, Ohio.

5 SIZES—
wood
or wire
handles



THE OHIO BOXBOARD CO.

PLANTS: Rittman, O. • Youngstown, O. • Cuyahoga Falls, O.

Pittsburgh, Pa. • Fairbanks Containers, Inc., Middletown, O.


Western Containers, Inc., Lockport, N. Y. • Champion Containers, Inc., Plymouth, Mich.

ARE YOU A SEED SALESMAN?

Add to your income by selling AMERICAN VEGETABLE GROWER. It's a logical combination for AMERICAN VEGETABLE GROWER will help your customers get the best results from the seed you sell them. Make every call pay! This means additional cash for you regardless of whether you sell a seed order or a subscription to AMERICAN VEGETABLE GROWER.

Write today for our liberal, nursery agents' plan. Address:

EDWARD MEISTER, General Manager
AMERICAN VEGETABLE GROWER
WILLOUGHBY, OHIO




Make your own

FLAKED ICE

for as little as

\$1.50 to \$2.00 per ton

**BURGE
FLAKED
ICE
MACHINE**



A fully automatic Ice Machine—Ice starts immediately. It's economical—you pay only for the ice you use. Occupies less space than the storage bin. Large capacity—3 to 12 tons per day. Installation is simple—can be connected to present refrigeration system or we can furnish complete with compressor, etc. Use coupon or write.

BURGE ICE MACHINE COMPANY

Dept. A, 654 W. Washington Blvd. Chicago 6, Illinois

Branches: Ft. Wayne, Ind.; Green Bay, Wis.; Knoxville, Tenn.

BURGE ICE MACHINE CO., Dept. A,
654 W. Washington Blvd., Chicago 6, Ill.
Please send bulletin on BERGE Flaked Ice Machine.

Name _____
Company _____
Address _____
City _____
Zone _____
State _____

Steaming Is OLD HAT!



Make Old Soil Like **NEW**

with **Larvacide**
EFFECTIVE • EASY • ECONOMICAL
SOIL FUMIGANT

Read what Dr. A. W. Dimock, Plant Pathologist at Cornell University, wrote in the N.Y. State Flower Growers' Bulletin #107, July 1954:

"With reasonably tight bottoms and sides of the benches, and with covers of pure plastic or plastic impregnated sheeting, it should be possible to attain as good sterilization and weed control with chlorpicrin as with steam."

**INCREASE YIELDS—CONTROL MOST SOIL-BORNE DISEASES,
"DAMPING-OFF, WILTS, ROOT ROTS," NEMATODES AND
MANY WEED SEEDS WITH SINGLE APPLICATION**



**For Best Results, Use Plastic Soil Covers—
For best plastic cover, use LARACOVER**



Write for FREE Technical Literature and Prices

Larvacide Products, Inc.

117 Liberty St.
New York 6, N. Y.

1515 Third St.
San Francisco, Calif.

**MORE IRRIGATION
AT LOWER COST
with the New PIR HALE PUMP**



**Compact—Rugged—Flexible
WILL PUMP** 500 GPM at 100 PSI
600 GPM at 85 PSI
700 GPM at 75 PSI

Write today for literature to Dept. AY
State No. of Acres and Water Source

HALE FIRE PUMP COMPANY
Conshohocken, Pa.

WHAT'LL IT MIX WITH?

1956 completely revised chart

That is the important question in economical spraying. The compatible nature of spray materials is mighty important in safe and effective spraying and AMERICAN VEGETABLE GROWER has produced an ingenious

SPRAY COMPATIBILITY CHART

which tells at a glance just what materials will mix safely. Printed in three colors, mounted on Bristol board paper, it is an accurate guide in the safe and successful mixing of all spray chemicals.

Better send for a copy—25¢ each
To foreign countries single copies—75¢ each

Quantity Prices on Request

AMERICAN VEGETABLE GROWER
Willoughby, Ohio

Enclosed is 25c. Send Compatibility Chart to:

Name _____

Address _____

City _____ State _____


FIVE DIFFERENT WAYS

(Continued from page 12)

grow a variety of vegetables for their roadside market as well as for wholesale. They place quality at the top of their produce requirements, and pest control is a big factor in attaining this.

Muskmelons, one of their prize crops, get a rotenone dust early for control of beetles. Later, when vines begin to run, a spray mixture of parathion and Zerlate is used for control of beetles, blight, anthracnose, and other pests.

Control of tomato and sweet corn pests on the Adams' Fairacre Farms is similar to the Henrys', with a few differences. They begin spraying very early for control of aphids which spread the Stewart's wilt bacteria. For aphid control they use a DDT emulsion applied with a weed sprayer, first just as the corn breaks through



Cabbage with severe damage from cabbage looper, a pest often defying control. Endrin and dieldrin are giving good control in some areas.

the ground. Two more applications at five-day intervals are made.

Early tomatoes at Fairacre Farms receive no sprays, as harvest is over before insects and diseases become destructive. Later varieties receive applications of fixed copper and Zerlate. Materials are alternated, with Zerlate in the first spray, copper in the next. Applications are repeated every week or 10 days until four sprays have been applied. This program keeps blights and anthracnose to a minimum.

The Adamses find their Hardie Aero Mist sprayer quite satisfactory for control and apply materials up to 6X concentration. With sweet corn they can cover eight rows at a time, doing a good job much faster than formerly with a boom sprayer.

The Blodgett Way

Stephen W. Blodgett grows 1500 acres of Black Valentine green beans in the lower Hudson Valley near Fishkill, N.Y. Leafhoppers, aphids, and other leaf-feeding insects are

AMERICAN VEGETABLE GROWER

major p
tory wit
parathion
every five
and even
hot, as le

The Sea

At Se
Farming
dreds of
for their
many oth
plication
and dis
50,000 a
effectiver
Seabrook
equipment

The d
is the m
individu
a swath
of the l
Iron Age

Pea a
peas, and
one airp
thion du
The rem
plication
fast bui
the peas
Root rot
some dis
and is c

Down
beans in
Farms c
ing meth
Hyre of
Newark,
mines th
which is
forecast.
sprays a
has kept
year Se
to one o
rials, for
Insect
many,
Mexican
cently th
are easi
spray ap
and mite
son a ne
tried for
plication
control c

The Ge

Al G
Ohio, gr
on his 90
land mar
duced hi
didn't. C
pays str
schedule

(C

MAY, 195


WAYS

(12)

s for their
for whole-
at the top
ents, and
in attain-

their prize
early for
when vines
e of para-
or control
nose, and

sweet corn
re Farms
with a few
aying very
ds which
bacteria.
e a DDT
d sprayer,
s through



om cabbage
Endrin and
some areas.

applications
de.
re Farms
st is over
s become
eceive ap-
and Zer-
ted, with
copper in
repeated
ntil four
This pro-
thracnose

r Hardie
atisfactory
ials up to
sweet corn
at a time,
ster than
ayer.

ows 1500
een beans
ley near
s, aphids,
sects are
E GROWER

major problems. Control is satisfac-
tory with dust applications of 1%
parathion. Applications are made
every five days during wet periods
and every seven days when dry and
hot, as long as insects are a problem.

The Seabrook Way

At Seabrook, N.J., the Seabrook
Farming Corporation grows hun-
dreds of acres of peas and lima beans
for their freezing plant, as well as
many other vegetables. Airplane ap-
plication is frequently used in insect
and disease control to cover the
50,000 acres quickly. However, for
effectiveness of control the folks at
Seabrook prefer ground application
equipment on most crops.

The duster now in use at Seabrook
is the new Hardie model without
individual nozzle tubes which covers
a swath 32 feet wide. Sprayers are
of the boom type made by Oliver
Iron Age.

Pea aphid is the major pest on
peas, and is controlled very well with
one airplane application of 1% para-
thion dust on about half the acreage.
The remainder requires a second ap-
plication, mostly because of a very
fast build-up of aphids, as soon as
the peas break through the ground.
Root rot has been the only trouble-
some disease in Seabrook pea fields,
and is controlled by crop rotations.

Downy mildew is a problem on lima
beans in the East, and at Seabrook
Farms control begins with a forecast-
ing method developed by Dr. R. A.
Hyre of the USDA, now working at
Newark, Del. This method deter-
mines the time to begin spraying,
which is eight days after mildew is
forecast. Applications of copper
sprays at seven- to 10-day intervals
has kept mildew to a minimum. This
year Seabrook growers may switch
to one of the newer carbamate mate-
rials, for a less costly program.

Insects attacking lima beans are
many, but the chief ones are
Mexican bean beetle, worms, and re-
cently the two-spotted mite. Beetles
are easily controlled with dust or
spray applications of 1% parathion,
and mites to some extent. This sea-
son a new material, Kelthane, is being
tried for control of mites. DDT ap-
plications are the most economical for
control of worms in lima beans.

The Gerhart Way

Al Gerhart, at North Ridgeville,
Ohio, grows a variety of vegetables
on his 90 acres, mostly for the Cleve-
land market. Many times he has pro-
duced high quality crops when others
didn't. One reason is that Gerhart
pays strict attention to pest control
schedules, and follows what he terms
(Continued on page 30)

MAY, 1956



ROW CROP ATTACHMENT!

Now you can convert your high pressure row crop
sprayer to a 2-fan air blast unit for concentrate or
dilute spray application. This new Hardie 2-fan Jetaire
Row Crop Attachment gives you the same unmatched
performance in Row Crop Pest Control as the famous
Hardie Jetaire 2-fan Air Blast Row Crop Sprayer. Ask
your Hardie dealer to show you this wonderful new
unit. Write for catalog.

COMES COMPLETE

- Equipped with 2, 26 in. axial fans.
- The reversible housing can be rotated through 220 degrees
by tractor seat controls—sprays on either side.
- Adjustable vanes provide easy direction control.
- Delivered ready to run with everything needed for in-
stallation on sprayers with either wood or steel tanks.
- Easily dismounted when not in use. Attractive low price.



The Hardie Jetaire is the only 2-fan air
blast row crop sprayer. Its performance
in big volume row crop pest control has
established unmistakably that 2 fans
are better than one in a row crop sprayer,
as well as in fruit tree sprayers. Leading
row crop growers all over America are
finding new ease, speed, economy and
efficiency in pest control with the 2-fan
Jetaire. Write for complete data.

Air Blast and High Pressure Sprayers,
Dusters, Irrigation Systems.

HARDIE

PEST CONTROL EQUIPMENT

SOLD AND SERVICED

Ask your dealer.

The Hardie Manufacturing Company, Hudson, Michigan
3825 Santa Fe Ave., Los Angeles 58, California
1435 N. W. Northrup St., Portland 9, Oregon
In Canada, C. W. Lewis & Son, Ltd., Grimsby, Ont.
Export Dept., Book Tower, Detroit 26, Michigan

Please send Catalog.

Name _____
Address _____
City _____ State _____

GREAT LAKES 659-G

New, improved Great Lakes lettuce, has dark green color with solid well-protected heads. This lettuce is heat, tipburn, and frost resistant and is a very heavy yielding lettuce that has proven itself from Maine to California in hot weather.



One of our 20 lines of Great Lakes lettuces.

ASK YOUR SEED DEALER FOR DESSERT'S SEED

WE ARE PRIMARY GROWERS OF ALL VEGETABLE SEED
SPECIALIZING IN QUALITY AND NEWNESS OF VARIETIES
AND GROW ALL LINES OF HYBRIDS

MAIN OFFICE

DESSERT SEED CO., INC.


P. O. Box 181

El Centro, California

MAKE SPACE RESERVATIONS NOW
FOR IMPORTANT JULY DIRECTORY ISSUE

The Newest Method in Weed Control THE BUDDING INROW WEEDER

Cut Your Weeding Cost



Growers who are using this weeder claim it to be the best ever, for cutting weeding cost.

One unit will do the work equal to hand labor of eight to ten men, practically eliminating all hoeing or hand weeding. Can be used in celery, cabbage, sweet corn, beans, strawberries, set onions, asparagus, mint, peanuts, gladiolus and other planted crops.

Works excellently on bedded crops and can also be set up for multiple row work. Cultivates around each plant at speeds of 4 to 6 miles per hour without injury to the crop.

Write for complete information today.

BUDDING INROW WEEDER CO.
Caledonia, Michigan

This Year Give Your
Seeds and Cuttings
A Real Break
With **SCOTTY** —



This Scotty 15 ft. by 35 ft. cash price,
only \$982.00.

**The LOW COST Greenhouse
That Goes Up Fast**

Would you say that propagation is the most important stage of plant culture? It's a debatable question—But one thing you can be sure of is that seeds and cuttings propagated in a Scotty greenhouse are getting the best start possible and under conditions of temperature and humidity that you control—not mother nature.



**Set up Scotty
yourself.**

Completely prefabricated parts are simply assembled into foundation spanning frames as shown at left. Assembly progresses rapidly.

**Write for
Free Catalog**

LORD & BURNHAM
IRVINGTON, N. Y. • DES PLAINES, ILL.

FIVE DIFFERENT WAYS

(Continued from page 29)

a "preventive program." His dusting schedule often begins before any signs of the pest appear.

Cucumbers are planted early in the spring when weather is often too wet for any field work after planting. So Al rigged a duster on his small crawler tractor and dusts regularly from the time plants break through the soil. He keeps a dust on all the time, making applications after every rain. Sometimes it means dusting every day. Last year when mosaic knocked out many neighboring fields, Gerhart harvested one of his finest and largest crops of cukes.

First dusts on cucumbers are with $\frac{3}{4}\%$ rotenone. When the second leaf develops Gerhart switches to a mixture of rotenone and Crag 658, continuing this mixture until about harvesttime. If dusts are necessary near



Muskmelon on farm of Ralph Adams and Sons, Poughkeepsie, N.Y., are on a rotenone-parathion-Zerlate program for insect and disease control.

or during harvest, rotenone alone is used. If mildew becomes a problem in the fall, a mixture of rotenone and 10% Zerlate is used.

Squash get about the same treatment as cucumbers as to frequencies of dust applications. The material is usually rotenone, although purified DDT is sometimes used. Applications are made after every rain, or about every five or six days if no rains occur. Mildex proves a pretty good control for mildew in the fall.

Cabbage and cauliflower are major crops on the Gerhart farms and often present the biggest insect control problems. A dust program begins a week after plants are set in the field and continues at 10-day intervals. Worms and loopers are controlled with a dust mixture of 3% DDT and 1% parathion used at the rate of 35 pounds per acre. If dusts are required late, within 30 days of harvest, then rotenone alone is applied.

This program has failed to control cabbage looper in recent years however. Last year Gerhart tried endrin for control of this pest without much success, then made one application of

(Continued on page 31)

AMERICAN VEGETABLE GROWER

tin Pari
growing
Parish
frozen
broccoli
grown c
market
however
is April
tin, Poi
St. Jam
areas fo

Tomato
most of
north L
shifting
Louisian
mines a
industry
The crop
stores an

Irish
have bee
a long p
stimulate
crop in t
a shift t
and new
Louisian
strong in
the early
in a pos
newer, h
the Red
introduce
ard vari
to a half
on a forw

Sweet
years L
all other
and mar
the intro
variety a
the Unit
of the ol
copper sh
flesh, it
to mainta

Today
per cent
grown in
fourteen
are shipp
and toda
duction i
are aroun
shippers

The s
grown as
to supply
foundation
Rico. Sin
ana stati
eration w
sweetpot
the USD
potato to
important
introduce

WAYS

(29)

is dusting
fore any

arly in the
en too wet
nting. So
his small
regularly
k through
on all the
fter every
s dusting
n mosaic
ing fields,
his finest

s are with
econd leaf
to a mix-
658, con-
about har-
ssary near



ns and Soss,
ne-parathio-
ease control.

e alone is
a problem
enone and

me treat-
requencies
material is
a purified
applications
or about
no rains
pretty good
all.

are major
and often
et control
begins a
n the field
intervals.
controlled
DDT and
rate of 35
s are re-
f harvest,
ed.

to control
ears how-
endrin
out much
ication of
(31)

E GROWER

LOUISIANA

(Continued from page 11)

tin Parishes. This industry is also growing rapidly in Avoyelles Parish. A considerable amount of broccoli is now being grown for the frozen food industry. Locally-grown cabbage can be found on the market from October through June, however, the main shipping month is April. The parishes of St. Martin, Pointe Coupee, Ascension, and St. James comprise the principal areas for this crop.

Tomato: Until a few years ago most of the tomatoes were grown in north Louisiana, but the industry is shifting to several of the south Louisiana parishes, mainly Plaquemines and St. Bernard, and this industry is growing rather rapidly. The crop is being sold to large chain stores and buyers.

Irish Potato: While Irish potatoes have been grown to some extent for a long period of time, World War I stimulated the production of this crop in this state. However, due to a shift to western production areas and new methods of packing, the Louisiana industry has not been as strong in recent years as it was in the early 30's. It is now, however, in a position to go forward since newer, high-yielding varieties like the Red LaSoda have been bred and introduced. It outyields the standard variety, Triumph, by a third to a half, and this industry is now on a forward move.

Sweetpotatoes: For the past 30 years Louisiana has led practically all other states in the production and marketing of this crop. With the introduction of the Porto Rico variety and with the introduction of the Unit I Porto Rico, a mutation of the old type, which has a better copper skin color and a dark orange flesh, it was possible for Louisiana to maintain its leading position.

Today Louisiana ships around 70 per cent of the commercial crop grown in the United States. Ten to fourteen thousand car equivalents are shipped from this state annually, and today almost half of the production is canned or frozen. There are around 25 canners and over 60 shippers in the state.

The sweetpotato industry has grown as the result of being able to supply the growers with the best foundation seed of the Unit I Porto Rico. Since 1937, when the Louisiana station, working in close co-operation with most of the important sweetpotato producing states and the USDA, first induced the sweetpotato to bloom and set seed, many important new varieties have been introduced which are resistant to

diseases and which have a higher nutritional value than some of the standard varieties. While the impact of these new varieties has not been fully felt, the industry is rapidly growing, not only in this state, but in other areas. In Louisiana the principal production areas are St. Landry, Lafayette, Evangeline, Arcadia, West Feliciana and West Carroll Parishes.

The Louisiana station maintains a Sweet Potato Research Center which is devoted entirely to the improvement of this crop through breeding and production of foundation seed. This center is located in Franklin Parish at Chase. The Louisiana station not only produces seed for its own breeding program but also furnishes many other co-operators with seed potatoes and true seed.

THE END.

FIVE DIFFERENT WAYS

(Continued from page 30)

dieldrin dust in August about a month before heading, with good control. Other growers have reported control with endrin.

Control of cabbage maggot on the Gerhart farm is with a 5% dieldrin dust applied directly over the row just after planting and repeated in six days.

Peppers get a dust application of Zerlate and parathion two or three times before bloom. This program has held insects to a minimum.

About the time tomato plants begin to bush up, perhaps three or four weeks after setting, they get their first dust application. This is a mixture of Manzate and Rhothane at the rate of 30 to 35 pounds per acre. This is followed in a week with Manzate alone. In another week this is repeated. A week later the fourth application is made with the Manzate-Rhothane mixture.

These four dusts are the basic control program for early and late blights and anthracnose. If the latter disease persists, a dust of 10% Zerlate is made just before ripening. Weather has much to do with the number of late dust applications.

This year Gerhart is changing his dust program on sweet corn from a straight 10% DDT to an alternating program. The first three dusts are with a 5% DDT, the fourth with a 2% parathion, and the fifth with a 10% DDT. This is based on a schedule of five applications beginning just when the first corn borer eggs hatch and repeating every five days. About 40 pounds of dust are applied per acre.

THE END.

WADE RAIN Leads the way

TO BIGGER YIELDS
... EARLIER HARVESTS
... BETTER QUALITY



Exclusive Features
SAVE TIME
and LABOR!

WADE RAIN
Sprinkler Irrigation



Features that make Profit!

WADE RAIN GIVES YOU ALL THE FAST, positive lateral moving advantages — as important to row crop growers! Why not have all the benefits of WADE RAIN Sprinkler Irrigation?

WADE RAIN SPEEDS PLANT GROWTH! Because, in every row, the early, better growth, means bigger yields and increased profits.



FREE PLANNING FORM
"FARM-FACTS" makes it easy to plan... Use Coupon

3 Revolutionary MECHANIZED LATERAL MOVES including POWER-SHIFT. Moves 1/4-Mile Lines as Unit. Tractor stays out of Crops!

ASK WADE RAIN Dealers About 5-YEAR PMA PURCHASE PLAN

SINCE 1865
Hqrs.—1919 N. W. Thurman St.
PORTLAND 9, OREGON
COLUMBUS 15, OHIO SEATTLE 4, WASH.
343 North Front St. 532 First Ave., South

Send me free "Farm-Facts" Planning Form and illustrated Wade Rain Literature.

NAME.....

ADDRESS.....

CITY..... STATE..... W-725

MAY, 1956



Uses include: Building soil humus by plowing down with cover crops or crop residues... fertilization... weed control.

Free Booklet: "AERO Cyanamid, what it is and how to use it." Write to American Cyanamid Company, 30 Rockefeller Plaza, New York 20, N. Y., or see your dealer.

SENSATIONAL!! New Low Cost Plastic Greenhouse Heating

WITH **KUR-MOR**

(USING L-P OR BOTTLED GAS)


50,000 BTU Input
Rugged Construction
No Vent Required

100% Safety Pilot
Directional Heat
Flow
Manual or Automatic Control

is Being Used With Outstanding Results

WRITE **PAT. PEND.**
BLUE GRASS BUTANE CO., Inc.
2417 Nicholasville Pike Lexington, Ky.

"ZON" SCARECROW



Imported from Holland • Portable • only 22 lbs.
Rugged construction. Safe, simple. No pilot light... years of proven economical service. Gives positive protection. Operates on carbide or acetylene tank for many days' operation without refueling. Retail price: \$50.00. Immediate delivery. For name of nearest dealer, write:

B. M. LAWRENCE & CO.
255 California St., San Francisco 11
Phone EXbrook 2-3310

AIR-BLAST SPRAYING

(Continued from page 9)

of sprayer designs have been carried on at the Ohio Experiment Station, in co-operation with the USDA Engineering Laboratory at Toledo, during the past five years. In 1952 a large air-blast sprayer capable of delivering 26,000 cubic feet of air a minute at an outlet velocity of 90 miles per hour was compared with the usual type of hydraulic sprayer equipped with an off-side boom in several different tomato fields. Foliage and fruit diseases were comparatively scarce.

The control obtained with the two machines was approximately equal when a 2X formulation of maneb was applied at 80 gallons per acre. When the air-blast machines were used to apply 2X, 2.7X, 4X, and 8X formulations at 80, 60, 40, and 20 gallons per acre, respectively, the degree of disease control obtained with the three higher rates was very similar, but 20 gallons of an 8X concentrate did not do as well.

Comparative Results

At Fremont, Ohio, in 1954, an experimental air-blast sprayer was used to apply maneb to tomatoes in 4X, 5.3X, 8X, and 16X formulations at 40, 30, 20, and 10 gallons per acre. At 40 gallons per acre the air-blast application gave slightly better control of fruit rots than was obtained with a hydraulic sprayer. It happened that the 30-gallon per acre rate (using a 5.3X concentrate) gave better control than 40 gallons, or than 20 and 10. The reason for the better performance at 30 gallons was not apparent, but it did show that good disease control could be obtained using an air-blast sprayer with comparatively low-gallonage applications.

In another comparison between these two types of spray applications made at Apple Creek, Ohio, in 1953, the hydraulic machine gave slightly better average control of a rather heavy infection of anthracnose fruit rot of tomato than did the air-blast sprayer when both machines were used to apply maneb at 20, 40, and 80 gallons per acre.

This situation was reversed at Toledo in 1954 in a field of late tomatoes where early blight was scarce and anthracnose rot was of only medium severity. In that experiment the air-blast application gave the better control of anthracnose at comparable gallonages.

In still another experiment of a similar nature at Bloomdale in 1955, where at least one-third of the fruits in the untreated check plot were infected with anthracnose, two different air-blast sprayers gave considerably better control of the disease than was

obtained by spraying with a hydraulic boom.

The data obtained in these experiments, conducted over a period of four years, indicate that, at comparable gallonages, the air-blast sprayer may be expected to give as good control of most of the foliage and fruit disease of tomato as can be obtained with hydraulic equipment, provided weather conditions are reasonably favorable for air-blast spraying.

Width of Swath

To get the best results with air-blast equipment the grower should avoid using a single swath much wider than 40 feet (80 feet double).

In 1953 a detailed study was made of the decrease in air velocity at increasingly greater distances from the air outlet of a comparatively large air-blast sprayer especially designed for treating row crops. The machine had a capacity of about 26,000 cubic feet of air delivered at approximately 90 miles per hour at the point where the nozzle manifold was mounted.

The average velocity during trials conducted on five different days at a point of 6 feet from the air outlet was 23.7 miles per hour. It was nearly as great at 18 feet with a value of 18.8 m.p.h. At 30 feet it was 15.5 miles. However, at 42 feet from the point of air emission the rate of air movement had decreased to 8 miles per hour, with a still further drop to 5.3 miles at 54 feet. The average velocities at the 6-, 18-, and 30-foot stations were about 4 miles per hour greater when the air-blast was traveling with the wind than when it was moving against it.

This drop in the rate of air movement with an increase in distance from the air outlet caused a considerable decrease in the quantity of spray material deposited on the foliage. The amount of copper per square centimeter of leaf area was 21, 17, 14, 6, and 5 micrograms at points 6, 18, 30, 42, and 54 feet from the sprayer, respectively.

These data indicate that the ability of a machine of this type, which was similar in size and design to many of those now in use, to carry and distribute spray material in a satisfactory manner decreases rapidly beyond a distance of 30 feet, until at 42 feet the spray deposit may be only one-third of that at 18 feet. This quantity (6 micrograms per square centimeter) would, of course, be doubled as the sprayer returned in going the opposite direction down the field, assuming that it was not over 42 feet away as it passed any given point along the swath.

AMERICAN VEGETABLE GROWER

How t
any other
crease in
is illustra
tained in
that insta
copper in
each squa
in the cer
15, comp
grams fro
the edges
mum valu
with a hy
15 microg

This v
across the
to be a
sprayer, b
that the m
per unit o
large as
sprayer. A
copper fro
blast swat
that to be
same widt
sprayer.

The rea
not clear,
of the spr
by the foli
across the
falls to th
ters a lea

IN
YOUR
BAND
Op
RUBB
ALLI
ALLIANCE, C

How the quantity of copper (or any other spray material) may decrease in the center of a double swath is illustrated by the deposit data obtained in the Fremont experiment. In that instance the average quantity of copper in micrograms recovered from each square centimeter of leaf surface in the center of the swath was only 15, compared with 25 to 30 micrograms from leaves only 12 feet from the edges. The maximum and minimum values of a similar swath treated with a hydraulic sprayer were 19 and 15 micrograms, respectively.

This variation in deposit values across the air-blast swath might seem to be a serious indictment of the sprayer, but it is interesting to note that the minimum of 15 micrograms per unit of leaf surface was still as large as that for the hydraulic sprayer. Also, the total recoverable copper from the foliage of an air-blast swath is often 50% greater than that to be found on a swath of the same width treated with a hydraulic sprayer.

The reason for this difference is not clear, but it seems likely that more of the spray material is intercepted by the foliage as it travels horizontally across the tops of the plants, and less falls to the ground before it encounters a leaf than when it is forced

vertically downward from the nozzle of a boom mounted on a hydraulic sprayer.

The experimental data so far obtained indicate that most air-blast sprayers, when properly used, are capable of giving a sufficiently good degree of disease control on such row crops as potato and tomato, and thus probably on many other crops also, that their use for that purpose may be recommended.

Specific Recommendations

However, certain specifications must be met by both the sprayer and its operator if best results are to be obtained.

1) A sprayer of average size should have the capacity to deliver at least 15,000, and better still 17,000 or more, cubic feet of air per minute at an outlet velocity of 80 to 100 miles per hour.

2) The pump should have a capacity of at least 15 to 20 gallons per minute and be capable of developing pressures of 500 to 600 pounds per square inch (a pressure of 400 p.s.i. is most commonly used).

3) Single swath widths should seldom exceed 40 to 45 feet (80 to 90 feet between sprayer driveways) and the sprayer should not be operated when the rate of air movement


(wind) is in excess of 6 or 7 miles per hour.

4) It is possible that the recommended rate of application may some day be dropped to as little as 20 gallons per acre, but at present the use of 30 to 40 gallons seems most likely to give good results. The corresponding spray-mixture formulations for these gallonages are an 8X at 20, a 5.3X at 30, and a 4X at 40 gallons per acre.

Specially designed, or extra large, machines may exceed some of the specifications given above, but the operator should be very certain that he can spray wider swaths than those suggested, use his sprayer in higher winds, or use more concentrated formulations at lower gallonages than those suggested, before he takes a chance on a lesser degree of disease control.

Finally, the material presented in this article should not be interpreted as an unqualified endorsement of the air-blast sprayer, in its present stage of development, as the best means of controlling vegetable diseases. However, it now appears very likely that this type of equipment will gain steadily in popularity and may eventually replace present-day hydraulic sprayers with all but the small grower whose fields or plots may be too small for its use.

THE END.



Banding with *Colorful ARCO Open Ring*

RUBBER BANDS Sell **MORE** Vegetables —

Yes, **COLOR DOES SELL MORE VEGETABLES** and Bright colored ARCO Rubber Bands with the distinctive "Open Ring" feature makes vegetable banding a "snap"—You can do it with one hand!

NO MORE BRUISED OR DAMAGED VEGETABLES—Rubber Bands Hold firmly but Gently, don't damage tender vegetables—Do keep them neat, clean, and easy for the dealer to display attractively.

**INCREASE
YOUR VEGETABLE
SALES**
BAND 'EM WITH
Open Ring
RUBBER BANDS

WRITE TODAY
for Samples & Prices

ALLIANCE RUBBER COMPANY

ALLIANCE, OHIO • CHICAGO, ILLINOIS • HOT SPRINGS, ARKANSAS
69 W. Washington

ALLIANCE RUBBER COMPANY


69 W. Washington CHICAGO, ILLINOIS

FIRM NAME

ATTENTION

ADDRESS

AVG-5



The BOOK EVERY GROWER HAS WANTED

- PRACTICAL
- UNDERSTANDABLE
- UP-TO-DATE

Written in simple, non-technical language, the **Fruit Pest Handbook** tells you why, when, and how to control fruit pests. Written for the grower by AMERICAN FRUIT GROWER editorial experts. Well-illustrated, easy to read, this 100-page pocket-size book will make the most complicated pest control problem simple and easy.

Every recommendation has been carefully checked to insure results. Compiled from the pages of AMERICAN FRUIT GROWER during the past several years, the **Fruit Pest Handbook** will be your most important reference and guide.

Order your copies today. Use convenient coupon below—just attach your check or money order.

\$2.00
PER
COPY

Books sent
postpaid.

AMERICAN FRUIT GROWER
Willoughby, Ohio

Please rush _____ copies of Fruit Pest Handbook. My remittance is enclosed (check or money order).

Name _____

Address or RFD _____

Town _____ State _____

KILL 20 MILLION WEEDS

(Continued from page 13)

grower were: What per cent spray is best? Which stage of the onion will tolerate a 1%, 1½%, 2% solution? What time of day is best to spray? How large a weed will the 1%, 1½%, and 2% kill?

Precision chemical weed killing resulted after worry, failures, disappointments and burned onions. However, many weeds can now be controlled successfully in onions with potassium cyanate.

Seed onions will tolerate a 1½% cyanate spray (6 pounds in 48 gallons of water) in the crook or tight-fold stage. Once the flag is up (the 7 stage), they will be hurt. If the spray is too late, a hand weeding job is necessary.

When the first "true" onion leaf is 3 inches long, a 1% spray can be used. The difference between a weed which can and cannot be killed with a 1% solution is about three days. Cyanate doesn't kill large weeds.

As the onion becomes taller, the cyanate solution can be increased to 2% if the spray is directed and kept off the onion leaves. The soil and weeds should be sprayed, not the onions. Cyanate kills many soft, broad-leaved annual weeds except lamb's-quarters. Rarely will cyanate kill purslane that is larger than a dime.

Special grade cyanamid, 75 pounds per acre applied before seed onion emergence, works successfully on muck soils when soil moisture is good. Stoddard Solvent applied before onion emergence kills the few weeds that are up at that time.

Chloro IPC, which must have descended from heaven, kills purslane and chickweed. These two kinds of weeds have been called by many names too numerous or too uncouth to mention. CIPC, 2 quarts in 50 gallons of water per acre (on mineral soils), 4 quarts per acre (on muck soils) before seed onion emergence, gives excellent chickweed and purslane control. Later CIPC sprays can be applied as needed, at the same rates, to control these two weeds. Control lasts from two to six weeks. Good soil moisture is necessary for good results.

Set and Transplant Onions

Set onions will tolerate a 1½% cyanate when they are 3 inches tall. Transplants will tolerate the same about ten days after setting. Sprays should be directed to avoid the onion leaves. Later sprays of CIPC are used as needed (4 quarts per acre on muck soils; 2 quarts per

acre on mineral soils) for chickweed and purslane control.

Effective onion weed control is not simple. Growers should be cautious until they thoroughly understand the action of these chemicals on crops and weeds.

Peas: Premerge or Sinox PE, one quart in 30 to 50 gallons of water per acre when the peas are 2 to 4 inches tall, kills most of the broad-leaved weeds. Air temperature must be between 70° and 80° F. to kill weeds and avoid injury. Container directions should be followed carefully when sprays must be applied at higher or lower temperatures.

Spinach: Chickweed in the spring, fall, and winter, with purslane in the summer, keeps many a vegetable grower busy. Chloro IPC, at 2 quarts per acre in 50 gallons of water during the summer, will give control for four to six weeks. In spring and fall when air temperature is 60° F. or below, one quart is top limit per acre. **Important:** CIPC should be applied immediately after seeding; soil moisture must be good; CIPC should be used only on mineral soils for spinach weed control.

Sweet Corn: One-half pound of 2,4-D per acre applied when the corn is 2 to 4 inches tall, is an old standby. Later sprays usually affect the corn. Earlier sprays at this rate don't kill enough weeds since they haven't emerged.

Watermelons, Cucumbers, and Muskmelons: Alanap-1 applied soon after planting, at 2 pounds in 50 gallons of water per acre, has given control of many broad-leaved weeds and annual grasses. The effectiveness of this weed killer varies with soil type. Follow directions with extreme care, and go slow if it's your first time with this chemical.

Continue Tillage

Chemical weed control is a supplement for cultivation and not a replacement.

The action of the herbicides (weed killers) mentioned were observed on the respective crops on several hundred farms in central New York (Madison County—bean and onion country) and western Long Island (Nassau County and Staten Island) by the author over a period of seven years.

The action of weed killers varies from area to area, and even from farm to farm. Contact your local extension service for area recommendations. THE END.

AMERICAN VEGETABLE GROWER

Ans

Q

Don't let
Whether large
and stamp
AMERICAN V
Ohio.

CARROT BLA

I store my
to 3 feet wide
ground. The
long and I p
When I take t
and ready th
spots on the
Colorado

It sounds
caused by th
Black rot oc
drying prior
gressive inc
disease as t
35°F. To re
dle your ca
the temperat
possible and

TOMATO VAN

Where can
which was de
ment Station?

Seed of t
tained from
Associated S
St., New H
pee Co., Phi
Seed Co.,
Seed Co., C

HARDENED PI

What are t
proof cabbage

Nearly a
harden their
ever, there
plants, and
done in pla
will make
less, anythi
hardening t
tend to tou
transplantin
survival an
growth will
transplantin

Cabbage
temperature
by gradual
peratures.
State Univ
plants have
or 15° below
of time. H
hardening h
tures will
continued fo

MILDEW RESI

Where can
dew Resistant
in your article

Try Asso
St., New H
Box 778, L
Co., St. Lou
& Sons, Inc

ONION SET H

Where can
Colorado.

Try R. C
Hoover Rd.
Mfg. Co., 51
and Paramo
St., Stockto

MAY, 1956

Answering Your QUESTIONS

Don't let your questions go unanswered. Whether large or small, send them with a three-cent stamp for early reply to Questions Editor, AMERICAN VEGETABLE GROWER, Willoughby, Ohio.

CARROT BLACK ROT

I store my carrots in November in trenches 2 to 3 feet wide, 2 feet deep, and 2 1/2 feet above ground. The trenches are about 75 to 100 feet long and I put air vents about 12 feet apart. When I take the carrots out of the trench to wash and ready them for cello pack, they have black spots on them. How can I prevent this?—Colorado

It sounds as if the trouble is black rot caused by the fungus *Alternaria radicina*. Black rot occurs as a result of rough handling prior to storage. There is a progressive increase in the prevalence of the disease as temperatures increase above 32° 35°F. To remedy the difficulty, try to handle your carrots more carefully and keep the temperature in the trenches as low as possible and still avoid freezing injury.

TOMATO VARIETY

Where can I obtain seed of the Kopia tomato which was developed at the Mississippi Experiment Station?—New York.

Seed of the Kopia tomato can be obtained from the following companies: Associated Seed Growers, Inc., 205 Church St., New Haven 2, Conn.; W. Atlee Burpee Co., Philadelphia 32, Pa.; Ferry-Morse Seed Co., Detroit, Mich.; and Service Seed Co., Crystal Springs, Miss.

HARDENED PLANTS

What are the advantages of hardened, frost-proof cabbage and tomato plants?—Minnesota.

Nearly all northern plant growers harden their plants before sale. However, there are no frost-proof tomato plants, and there is nothing that can be done in plant production practices that will make them frost-proof. Nevertheless, anything that is done in the way of hardening that will slow up growth and tend to toughen tomato plants prior to transplanting, improves the percentage of survival and the rapidity with which growth will be resumed in the field after transplanting.

Cabbage plants will withstand cold temperatures if they have been hardened by gradually subjecting them to low temperatures. S. H. Wittwer of Michigan State University reports that cabbage plants have been known to withstand 10° or 15° below freezing for a short period of time. However, an excess amount of hardening by exposure to cold temperatures will likely induce bolting when continued for several weeks.

MILDEW RESISTANT CANTALOUPE

Where can I obtain seed of the Powdery Mildew Resistant strain No. 45 cantaloup mentioned in your article on Arizona?—New Jersey.


Try Associated Seed Co., 205 Church St., New Haven 2, Conn.; Ferry-Morse, Box 778, Detroit, Mich.; Corneli Seed Co., St. Louis 2, Mo.; and F. H. Woodruff & Sons, Inc., Milford, Conn.

ONION SET HARVESTERS

Where can I obtain an onion set harvester?—Colorado.

Try R. G. Bruner Mfg. Co., 22516 Hoover Rd., Van Dyke, Mich.; Lauridsen Mfg. Co., 514 N. 11th Ave., Greeley, Colo.; and Paramount Mfg. Co., 1615 E. Main St., Stockton, Calif.

MAY, 1956



LANDPLANE . . . the way to Increased Crop Yield & Lower Irrigation Cost!

Increase your profit with a Marvin Multi-Purpose LANDPLANE! Whether one acre or 1000, LANDPLANE does "correction leveling" better, cheaper and more efficiently. Smooth, level fields get even moisture penetration, uniform crop growth. Irrigation costs reduced as much as 50%! Levels, cultivates, smooths and rolls in one operation!

There's a LANDPLANE to fit even the tightest farm budget. Be sure it's a LANDPLANE — the original scientific land leveler, built by Marvin. Write for FREE catalog.



VEGETABLE PLANTS

Cabbage, Onions, Tomatoes, Pepper, Collards, Broccoli, Brussels Sprouts, Cauliflower, Eggplants, and Sweet Potatoes.

Best strains of leading varieties. Shipments catering to market gardeners' demands. Free price list.

TEXAS PLANT FARMS
JACKSONVILLE, TEXAS

ELLIS QUALITY TRANSPLANTERS



"THE COMPLETE LINE"


- CONVENTIONAL • HORSE DRAWN
- TRACTOR DRAWN • TRACTOR MOUNTED (1 and 2-row)
- MECHANICAL (3 Point Mounted)

For More Information, Write

D. R. ELLIS MFG. CO.
Verona, Wisconsin

EYE GLASSES by MAIL As low as \$1.00

WRITE for FREE CATALOG with 14 LENS SAMPLE CARD
Thousands of Customers
Est. 1929



QUALITY READING-MAGNIFYING OR BIFOCAL GLASSES FOR FAR AND NEAR
ADVANCE SPECTACLE COMPANY, Inc.
537 S. Dearborn St., Dept. VG, Chicago 5, Illinois

Priced Right!

Engineered Right!

Serviced Right!

McDowell
COUPLING
Portable Irrigation
SYSTEMS



See your McDowell dealer—he's a good man to know better!

For portable irrigation that works better with less work and less upkeep, see your McDowell dealer. He offers the best coupling made . . . will engineer a complete system for your needs. . . and he's staffed to give fast, expert service.




Write for
Free Booklet!

McDOWELL MANUFACTURING CO.
Pittsburgh 9, Pa.

NEW FOR YOU

—to increase your profits

Cut Weeding Costs



A new weeder which does much of the back-breaking work automatically is now on the market. Developed by a vegetable grower, one unit will do work equal to the hand labor of eight or ten men, practically eliminating hand hoeing or weeding. The new unit has been used very successfully with celery, cabbage, sweetcorn, beans, strawberries, set onions, asparagus, and other vegetable crops. The machine works perfectly on bedded crops and can be set up to do multiple row work if needed. It cultivates around each plant at speeds of 4 to 6 miles per hour without injury to the crop. Write C. Buddingh, Buddingh In-Row Weeder Company, Caledonia, Mich., for further details.


Precision Seeder

Much has been said and written about precision seeding. There can be no doubt that in actual use the precision seeder will save growers like ourselves money not only by lowering seed costs, but by increasing production. The seeder pictured here is one of the best. It has passed every test with flying colors, and I can recommend it without reservation. You can get all the information you need on the new precision seeder by merely writing H. W. Laros at S. L. Allen & Co., 3419 N. Fifth St., Phila-

delphia 40, Pa. You'll find that the S. L. Allen seeder includes features not found in many other types of equipment.

Know Your Motor


It is very important that every vegetable grower know how many hours his tractor motor or sprayer motor has been operating. With this knowledge the grower can set up a program of preventative maintenance which will save money and increase the life and efficiency of his equipment. The direct-reading meter pictured above is a simple, easy way to do this. Extremely accurate, yet inexpensive, it is available in 6-volt, 12-volt, 24-volt, 32-volt, 64-volt and 110-volt models. Why not write John W. Hobbs Corporation, Sales Division, Springfield, Ill., for full details?



Hydrocooling




Vegetable growers everywhere are finding that hydrocooling increases quality and insures higher prices at the market. The hydrocooler pictured above is an improved model and handles sweet corn,



carrots, celery, asparagus, beans, radishes, peas, cauliflower, spinach, lettuce and onions with dispatch. Why not write Milt Kahn, Food Machinery & Chemical Corps., P.O. Box 1718, Lakeland, Fla., for your free booklet on hydrocooling and what it can mean to you.

Pruner Weeder



For the small grower who is looking for a good pruner which at the same time can be used as a weeder around his house, the Ristila Pruning and Weed Cane is ideal. The pruner will handle branches up to $\frac{3}{8}$ inch in diameter. As a pruner, the equipment works easily and effectively. I have used one and have found it ruggedly constructed and built to last. The cutting knives are taken out of the jaw of the unit, and quickly you have a weeder. It is ideal for weeding in small fruits and taking the old wood out of raspberries, blueberries, etc. Get the details on this new tool by writing John A. Overhouse, Overhouse Eng. & Mfg. Co., Box 489, East Lansing, Mich.

Important Catalog

I have just finished reading one of the best vegetable seed catalogs I have ever read. This booklet is comprehensive and covers all the phases of vegetable growing along with a detailed description of the various seeds available. This booklet will be sent to all readers of AMERICAN VEGETABLE GROWER free of charge. It has over 128 pages, and is something you should certainly have on your desk. Why not write the Corneli Seed Company, 101 Chouteau Ave., St. Louis 2, Mo., and ask Mr. Joseph Burger to send you their catalog #14?



AMERICAN VEGETABLE GROWER

ELE
'BETTE

TOMAT
Electric

George B
The p
of cinder
of the sid
glass sash
deep, the
There is 3
the bed c
with outle
sections.
controls t

Barker
inches of
thermosta
He estima
tricity wi
move the

The ma
the same
and you
haul it, an

Barker
the heat
the stuff
up to 125°

Electricity
more unif

Electric
the hotbed
Barker po
materials
manure b

He exp
of the ne
E. Toles.

B

The G
Cynthia W
ican Gard

Company,
When th
book of ga
appeared i
New York

garden bu
bug books

Since th
chemicals
that it beca
579-page v
contains al
chemicals,
mendation
color illust
numerous
a valuable

in growing
mentals, fl
Dr. We
pathologist
as a physi
years ago.
Doctor in
she sprays
from May
to lecture a

MAY, 1956

ELECTRIC HOTBED 'BETTER THAN MANURE'

TOMATO plants are grown in an electric hotbed on the farm of George Barker, Mechanicsville, Va. The permanent 12-sash bed made of cinder block is 38 x 6 feet. One of the sidewalls which supports the glass sashes on a slant is 17 inches deep, the other is 12 inches deep. There is 360 feet of heating cable in the bed crossing it every 7 inches, with outlets for plugging in different sections. A thermostat in the soil controls temperature.

Barker covers the cable with 6 inches of dirt, sows the seed, sets the thermostat at 65° F., and forgets it. He estimates that \$15 worth of electricity will heat a bed until time to move the plants to the coldframes.

The manure needed to accomplish the same purpose would cost \$12—and you have to find the manure, haul it, and put it down.

Barker said, "You can't regulate the heat with manure. Sometimes the stuff will run the temperature up to 125° F. and burn up the plants. Electricity makes them grow a lot more uniformly."

Electrical materials and labor for the hotbed came to about \$135, but Barker pointed out, "You save on materials for the bed. The regular manure bed is a lot deeper."

He expects to get 8,000 plants out of the new electric hotbed.—George E. Toles.

BOOK REVIEW

The Gardener's Bug Book by Cynthia Westcott. \$7.50. The American Garden Guild and Doubleday & Company, Inc.

When the first edition of this handbook of garden pests and their control appeared in 1946, it was hailed by the *New York Herald Tribune* as "the garden bug book to end all garden bug books."

Since then, says the author, new chemicals came on the market so fast that it became imperative to revise the 579-page volume. This second edition contains all the new pests, all the new chemicals, and all the new recommendations for control. With full-color illustrations of 102 pests, and numerous line drawings, the book is a valuable aid to the home gardener in growing vegetables, fruit, ornamentals, flowers, or just plain grass.

Dr. Westcott was the first plant pathologist to go into private practice as a physician to gardens some 25 years ago. Operating as The Plant Doctor in New Jersey and New York, she sprays over 50 gardens a week from May to November and manages to lecture around the country besides.

MEDICAL

LATEST METHODS TREATING PILES, Fistula, Colon-Stomach disorders. Write for free book. **McCLEARY CLINIC AND HOSPITAL**, E345 Elms Blvd., Excelsior Springs, Mo.

MISCELLANEOUS

PEAFOWL, PHEASANTS, GUINEAS, Bantams, Waterfowl, thirty varieties Pigeons. JOHN HASS, Bettendorf, Iowa.

MAILING LISTS—100,000 CAREFULLY SELECTED names in the Agricultural Field, Individual lists of Potato Growers, Tomato Growers, Corn Growers and many others. Ideal for firms wishing to contact leading growers and dealers. For details write: **MACFARLAND CO.**, Box 2, 8 Elm Street, Westfield, N. J.

NOW YOU CAN'T AFFORD NOT TO KNOW soil analysis—ph, nitrogen, phosphate, potash—\$3.50. Satisfaction guaranteed. **TRACE DETECTION LABS.**, 4920 Plum Drive, Houston 17, Texas.

PLANTS

"OZARK BEAUTY" AMAZING NEW EVER-bearing strawberry 20th Century and Redrich Cross. It's vigorous, healthy, productive and wonderful flavor. Photos and circular, write **WINNS BERRY FARMS**, Westfork, Ark.

SALESMEN WANTED

FARMERS, DEALERS, AGENTS — MAKE extra money. Demonstrate, take orders, new proven nationally advertised **Gro-Green Liquid Fertilizer and Nitrogen Nutrients**. Full-part time. Samples and demonstrating outfit **FREE**. **CAMPBELL CO.**, Rochelle 91, Illinois.

WANTED—HARDWARE SALESMEN IN Ohio and Western Pennsylvania who are calling on hardware stores and garden supply stores to sell a non-competing line on a straight commission. An excellent way to add to your income. All inquiries will be kept confidential. Write **BOX 104, Hardware Retailer**, Indianapolis 4, Indiana.

VEGETABLE PLANTS

SWEET POTATO PLANTS, GUARANTEED. Nancy Halls, Portoricans. 500, \$1.50; 1,000, \$2.50; 5,000, \$11.75. "Bunch" Portoricans. 200, \$1.25; 500, \$2.25; 1,000, \$4.00. Planting guide free. **SUNSHINE PLANT CO.**, Gleason, Tenn.

OPPORTUNITY ADS

Only 25c a word for one-time insertion. 20c a word for two-time insertion; 15c a word for four-time insertion—**CASH WITH ORDER**. Count each initial and whole number as one word. **ADDRESS AMERICAN VEGETABLE GROWER**, Willoughby, Ohio.

FOR SALE—EQUIPMENT AND SUPPLIES

FOR SALE—ASPARAGUS TRIMMER and conveyor, vegetable tyers. **MANHATTAN GARDENS**, Manhattan, Illinois.

4-ROW HARBISON-PAINE 'MILTON' PRECISION planter with fertilizer attachment. **GEO. DEVRIES FARMS**, Evergreen Park, Illinois.

SAWDUST MAKES GOOD FERTILIZER

You can apply 15 tons of sawdust per acre to build rich top-soil rapidly, and greatly increase nitrogen level, if you use the **Active** process. Cheaper, more permanent, and better soil condition than with chemical fertilizer. No compost-heap making. Apply now; anytime. Rush postcard for free bulletin. "How to Turn Waste into Gold."

THE ACTIVE PROCESS, Bridgeton 18, Ind.

FOR SALE—REAL ESTATE

IF YOU ARE AN EXPERIENCED VEGETABLE grower and want a greater return for your efforts, in both income and healthy living, there is a sunny future for you in South Florida. We have available some of the finest farmland in the east coast, west coast or Everglades sections. You, like many successful growers, might find the soils and climate of South Florida to your liking. Wire or write **HAROLD RABIN CO., INC.**, produce brokers and shippers, P. O. Box 7, Belle Glade, Florida.

IRRIGATION EQUIPMENT

DRAGON DISTRIBUTORS WANTED— Looking for a trouble free money making sprinkler irrigation line? We manufacture **DRAGON**, one of the best. Also, **DRAGON** injectors for sprinkler-fertilization. Inquire today. **DRAGON ENGINEERING COMPANY**, 626 McClary, Oakland 21, California.

OPPORTUNITY ADS

BUY, SELL AND TRADE—Readers and business firms will get top advertising value at low cost from **AMERICAN VEGETABLE GROWER** "Opportunity Ads." These classified ads are widely read, widely responded to by **AMERICAN VEGETABLE GROWER'S** high-income readers throughout America. Our classified advertising department will run your ad under any heading and will guarantee correct insertion.

RATES—For one insertion, 25c per word. For two insertions, 20c per word. Four insertions, 15c per word. Count each initial or whole number as one word. **CASH WITH ORDER.**

AMERICAN VEGETABLE GROWER reserves the right to reject or alter any copy which does not meet its standards. Ads will be inserted in first available issues unless preferred month is specified by advertiser. Closing date is first of month preceding publication.

USE THIS INSERTION FORM TODAY

Please insert my classified ad in your "Opportunity Ads" section as follows:

PLEASE _____

PRINT _____

OR _____

TYPE _____

MESSAGE _____

HERE _____

Place my ad under the following heading: _____

I enclose \$_____ for _____ words. Based on _____-time insertion rate of _____c per word.

1 (insertion 25c per word) 2 insertions, 20c; 4 or more insertions, 15c

AMERICAN VEGETABLE GROWER | **WILLOUGHBY, OHIO**

Action That Nets Growers a Profit

STRAWBERRY growers in southwest Michigan last year sold their second largest crop on record for the second highest average price in history. The highest price, received a few years ago, was in a year when few strawberries went to market. Peach growers in the same area experienced a better market and price last year than in any other section of the United States.

Why?

The growers themselves have the answer. They feel their good returns last year stem from the \$20,000 they spent in well-directed sales promotion. R. F. Bittner, district marketing agent at St. Joseph, will agree with them.

What is the story behind their high returns? About 1950 strawberry growers in Berrien, Van Buren, Allegan, and Cass counties organized the Southwest Michigan Strawberry Growers Association. Their purpose was to further the berry industry in their area and start some promotional work.

Two years ago the peach growers informally organized the Michigan Fruit Sponsors, primarily for the purpose of promoting the sale of their fruit. Having little money and no definite plans, not much was done until last year.

When District Marketing Agent Bittner discussed these two promotional programs with William Anderson, chairman of the Michigan State Agriculture Commission, they concluded the two ought to work together.

A meeting was arranged with the officers of both grower organizations; George McIntyre, director of the State Department of Agriculture; Mr. Anderson and Mr. Bittner; and the Benton Harbor Marketing Board.

The idea met with fine response and the Southwestern Michigan Fruit and Vegetable Promotion Committee was formed. The organization is so organized that if and when the need arises it can promote almost any horticultural crop grown in the area.

To help in financing the promotional efforts, the Benton Harbor Marketing Board agreed to contribute to the program 20 per cent of the revenue from each grower-load of produce sold through the big market.

Mr. Bittner is chairman of the promotion committee, and the various

interests are represented as follows: Strawberry growers—Otis Klett and John Babcock, both of Hartford. Michigan Fruit Sponsors—Earl Steimle, Sodus Fruit Exchange, Sodus, and Wallace Heuser, Hartford. Benton Harbor Market Board—Reid Sprague, Benton Harbor. Farm Bureau—Peter Ruddell, Berrien Springs, for strawberries; Harry Zepik, Watervliet, for vegetables.

During the 1955 season this group met frequently to work out details of the promotional efforts. They hired the Paxson Advertising Agency of Benton Harbor to help set up and carry out promotional programs. Last season promotional efforts were concentrated on strawberries and peaches, with tomatoes, cantaloupes, and cucumbers getting some attention. Promotional work consisted of radio and newspaper advertising, store streamers and letters to wholesalers, brokers, and produce men informing them, ahead of time, of the promotions planned on a given product.

The promotional work was carried on in five states—in such principal market centers as Chicago, Detroit, Cleveland, Columbus, Indianapolis, Fort Wayne, South Bend, and Milwaukee. The response was gratifying even though the group had only \$20,000 to do the job.

The growers raised \$10,000 and the Michigan Department of Agriculture, which matches dollar for dollar the money raised by a farm organization

VEGETABLE CONVENTION



"Yowl Yowl Yowl This is what I mean by surprise!"

for the promotion of its products, matched it with an equal amount.

This year the committee expects to have considerably more money available and plans a more complete job of promoting horticultural crops. The organization is sparked with the kind of vitality that brings positive results.

We Are the Key

AGRICULTURE today is endeavoring to win its place in this, the power age, by putting to use every available technological development.

Fundamentally, our achievements are of a personal nature. One person may conceive an idea, and may even put it into practice. The influences and effects of the idea, however, may change the lives of countless other persons. We need think only of the discovery of electricity by Benjamin Franklin, or of the invention of the electric light bulb by Thomas Edison.

These ideas came to mind as we read recently of the rising productivity of the American farmer, his ever-increasing efficiency. Over the past 15 years production per acre on our farms has increased one-fifth. An hour of a man's labor today produces twice as much produce as it did 15 years ago.

Greater use of the products of scientific research brings increasing dependence upon them and upon those who develop machines and chemicals and devise ways of using them. Thus, a chemist in a chemical manufacturing plant, a machinist in a tractor factory, a research worker in an agricultural experiment station are as responsible for changes in our spray programs, cultural practices, marketing systems as we, the growers, are. We are all interested in the same achievement; use of technological developments to raise the efficiency of production which in turn raises our standard of living.

We, the growers, are the key to this whole process. What if there were no incentive to make any changes? Industries would suffer. Our ability to purchase and use the products of technology is basic.

Coming Next Month


Irrigation Issue
featuring

- Guides to Efficient Irrigation
- Better Crops with Sprinklers
- What's New in Irrigation Equipment?
- Irrigation Problems in Florida

AMERICAN VEGETABLE GROWER

John BEAN POTATO HARVESTING

...the proven way to bigger potato profits



MODEL 26 HARVESTER IN USE WITH BULK LOADING

REDUCE YOUR LABOR COSTS 50%—OR MORE!

Growers using John Bean Potato Harvesters have cut their labor costs in half — and see potential cost reductions of up to 75%. They're comparing their previous methods of hand digging, picking and loading with John Bean mechanical indirect harvesting and bulk loading. With the somewhat

slower method of direct harvesting, labor costs are still reduced appreciably.

Profitable operation with John Bean Harvesters is not limited to the large growers — it's been proven that farms with 30 acres or more, or neighbors combining acreage, can profit substantially.

ELIMINATE TRANSIENT LABOR PROBLEMS

No longer need growers be concerned with the difficulties of obtaining, supervising and housing large crews. John Bean Harvesters have replaced 30 to 40 pickers and allowed growers to use only their regular force. The faster, easier operation helps keep good men, too. John Bean Harvesters

have independent power units that do not rely upon tractor PTO, so that apron speeds can be regulated easily to ground speeds for better teamwork and maximum output. Digger and bulkloader are hydraulically controlled.

PRODUCE A BETTER, CLEANER, MORE MARKETABLE CROP

John Bean Harvesters give you the most complete separation of potatoes from stones and clods of any equipment on the market. Many growers say they get far less damage from cuts and

bruises than with hand digging, because the potatoes are picked up and dropped fewer times. You gain closer control over the quality of your crop and better opportunity for top prices.

JOHN BEAN MODEL 26 POTATO HARVESTER

is the field-proven, heavy-duty equipment for one-row direct or two-row indirect harvesting. Has 26" elevating conveyor and 30" sorting table.

JOHN BEAN MODEL 60 POTATO HARVESTER

especially designed for two-row direct harvesting, or four-row indirect. Has 60" elevator and 36" sorting table.

See your John Bean dealer for details on this profit-making equipment or write for catalog



John BEAN

Lansing 4, Michigan San Jose 1, Calif.
Div. of Food Machinery and Chemical Corp.

Still the biggest money-maker:

ASGROW BLACK VALENTINE



But—for that “extra-dollar” quality grow the choice ASGROW strain



You can be sure it's
the ASGROW strain
if it comes in an
Asgrow bag

1. straighter, trimmer pods
2. attractive, uniform color
3. excellent shipping qualities



Order now for late season plantings
From your dealer, or direct from our nearest address

ASSOCIATED SEED GROWERS, Inc.

Breeders and growers of vegetable seeds since 1856

Atlanta 2 • Cambridge, N.Y. • Indianapolis 25 • Exmore, Va. • Memphis 2

Milford, Conn. • Oakland 4 • Salinas • San Antonio 11 • Vineland, N.J.

International division: Asgrow Export Corp., Milford, Conn.

Distributors for Florida: The Kilgore Seed Company, Plant City